

# International Journal of Psychosocial Rehabilitation

ISSN 1475-7192

*(Proudly Serving our Web Based, International Readers Since 1996)*



**A WEB BASED PEER REVIEWED PUBLICATION FOR MENTAL  
HEALTH PRACTITIONERS, CONSUMERS & APPLIED  
RESEARCHERS**

[www.psychosocial.com](http://www.psychosocial.com)



Copyright © 2011 JDA, LTD. All Rights Reserved.

### **Journal Information**

This peer reviewed Journal was created in 1996 by practitioners, mental health program managers and mental health consumers to provide international practitioners, scholars and consumers with a forum to publish and discuss their work in program development, evaluation research, policy innovations, and therapeutic practices that have been successful in their particular region and cultures. IJPR is not associated with any university or governmental institution, nor is it part of any ‘old boy’ or other professional network. It was created to provide information to an international readership about issues related to psychosocial rehabilitation and associated topics.

Articles on psychosocial interventions, psychopharmacotherapy, mental health primary care, institutional and community care innovations, decentralization, policy changes, community & regionally based systems, and program evaluation are given particular attention. However, all articles that relate to psychosocial rehabilitation will be considered.

We invite comment from all readers on any and all subjects published in this journal, including the journal format itself. Feel free to comment on the Bulletin Board as well.

#### ***Current Editorial Board***

Dr. Nira Hariyatie Hartani – Malaysia

Asst. Prof. Dr. Mohd Armi Abu Samah – Malaysia

Rathi Aarti, BPTH, MPTh, – Cardiorespiratory – India

editor@psychosocial.com

#### ***Associate Editors***

Ass. Prof. Dr. I Wayan Suryasa, Indonesia

### **Publishing Information**

The International Journal of Psychosocial Rehabilitation is published by ADG, S..A .(ADG) ADG is a private non-profit, limited corporation in Honduras and operating in a limited capacity in the United

Kingdom. All information in IJPR is provided for personal use. Reproduction for publication or other use is strictly prohibited. Written permission must be obtained ADG, S.A., to reprint any information contained within the Journal, either in part or its entirety .

submission@psychosocial.com

## For authors

---

Scope (<https://www.psychosocial.com/scope/>)

Track Your Paper (<https://www.psychosocial.com/track-your-paper/>)

Submit Online (<https://www.psychosocial.com/submit-online/>)

Editorial Overview (<https://www.psychosocial.com/editorial-overview/>)

Instructions for Authors (<https://www.psychosocial.com/instructions-for-authors/>)

Privacy & Cookie Policy (<https://www.psychosocial.com/privacy-cookie-policy/>)

Terms & Conditions (<https://www.psychosocial.com/terms-conditions-of-use/>)



## Quick Links

About Publisher (<https://www.psychosocial.com/about-publisher/>)

Copy Rights (<https://www.psychosocial.com/text-book/>)

## Contact Us

**editor@psychosocial.com** (<mailto:editor@psychosocial.com>)

**Hampstead Psychological Associates,**

Suite B19, 110 Gloucester Road,

London, NW1 8JA.

United Kingdom

Copyrights © 2022 SDA, LTD. All Rights Reserved.

Issue 7

**The relationship between self-esteem and intimate partner violence (IPV): The mediating role of resilience (https://www.psychosocial.com/article/PR270001/13822/)**

**Authors:** Samina Ashraf Chaddher\*, Roselisa Binti Ahmad

**DOI:** 10.37200/IJPR/V24I7/PR270001 (https://doi.org/10.37200/IJPR/V24I7/PR270001)

**Pages:** 1-9

**Keywords:** Intimate partner violence, Domestic violence, Self-esteem, Resilience

Abstract (https://www.psychosocial.com/article/PR270001/13822/)

Download (/register-login)

**INNOVATION AND PRODUCTION POTENTIAL OF MESOSYSTEMS: ASSESSMENT AND MODELING OF DEVELOPMENT PROSPECTS (https://www.psychosocial.com/article/PR270002/13854/)**

**Authors:** Alexey I. Kljushkin, Marina V. Shinkevich, Nina A. Ivanova , Olga V. Popova, Oksana N. Ryabchenko , Konstantin I. Brykin

**DOI:** 10.37200/IJPR/V24I7/PR270002 (https://doi.org/10.37200/IJPR/V24I7/PR270002)

**Pages:** 10-20

**Keywords:** innovative production potential, mesosystem, metallurgical industry, industrial production index, innovative activity, high-tech and high-tech production, technological innovations, modernization

Abstract (https://www.psychosocial.com/article/PR270002/13854/)

Download (/register-login)

**Organizational Sustainability Through the Application of Holistic Human Resources Development Model (HHRDM) in Multi-National Companies of Penang, Malaysia (https://www.psychosocial.com/article/PR270003/13860/)**

**Authors:** Mohd Anuar Arshad, Arshad Mahmood \*, Shahid Khan, , Kaleemullah, Adeel Ahmed, Muhammad Salman Shabbir

**DOI:** 10.37200/IJPR/V24I7/PR270003 (https://doi.org/10.37200/IJPR/V24I7/PR270003)

**Pages:** 21-28

**Keywords:** Organizational Sustainability, Holistic Human Resource Development, Spiritual Quotient

Abstract (<https://www.psychosocial.com/article/PR270003/13860/>)

Download (/register-login)

**Study of Self-Compassion, Depression, Anxiety and Stress in male and female teachers in Government College of Delhi and Uttar Pradesh in India (<https://www.psychosocial.com/article/PR270004/13867/>)**

**Authors:** Kamini C. Tanwar\*, Megha Garg

**DOI:** 10.37200/IJPR/V24I7/PR270004 (<https://doi.org/10.37200/IJPR/V24I7/PR270004>)

**Pages:** 29-43

**Keywords:** College teachers; Depression, Anxiety and Stress (DAS); Self-compassion

Abstract (<https://www.psychosocial.com/article/PR270004/13867/>)

Download (/register-login)

**An Assessment on Level of Stress among the Workers in Cotton Mill Industries in India (<https://www.psychosocial.com/article/PR270005/13884/>)**

**Authors:** K.M.Ashifa

**DOI:** 10.37200/IJPR/V24I7/PR270005 (<https://doi.org/10.37200/IJPR/V24I7/PR270005>)

**Pages:** 44-52

**Keywords:** Stress, Healthiness, Self-esteem, Attentiveness, Cotton mill, Worker.

Abstract (<https://www.psychosocial.com/article/PR270005/13884/>)

Download (/register-login)

**Investigating the Effect of Acupuncture on Facilitating Placenta and Fetus Removal in Difficult Labors (<https://www.psychosocial.com/article/PR270006/13900/>)**

**Authors:** Yousef Kolookhi, Mozghan Modoodi, Mina Ataei , Rafat Rezapour-Nasrabad , Mahshid Bazrafkan , Zahra Mehdizadeh Tourzani, Maryam Mirzaei \*, Gohar Hagpanah

**DOI:** 10.37200/IJPR/V24I7/PR270006 (<https://doi.org/10.37200/IJPR/V24I7/PR270006>)

**Pages:** 53-60

**Keywords:** acupuncture, placenta and fetus removal, vaginal delivery, Shahroud hospitals.

Abstract (<https://www.psychosocial.com/article/PR270006/13900/>)

Download (/register-login)

**Investigation of New Chromosomal Abnormalities Associated with Recurrent Abortion (<https://www.psychosocial.com/article/PR270007/13897/>)**

**Authors:** Mina Ataei, Zohreh Moslehi, Rafat Rezapour-Nasrabad, Mozghan Modoodi, Parisa Farahbakhsh, Samira Saeed Rad\*

**DOI:** 10.37200/IJPR/V24I7/PR270007 (<https://doi.org/10.37200/IJPR/V24I7/PR270007>)

**Pages:** 61-68

**Keywords:** recurrent abortion, genetic investigation, Chromosome abnormality, Karaj.

**Authors:** Amellia Mardhika, Agus Sulistyono, Emuliana Sulpat, Susilo Harianto

**DOI:** 10.37200/IJPR/V24I7/PR270726 (<https://doi.org/10.37200/IJPR/V24I7/PR2700726>)

**Pages:** 7576-7586

**Keywords:** mother, breastfeeding, counseling, exclusive, rct.

Abstract (<https://www.psychosocial.com/article/PR270726/18661/>)

Download (</register-login>)

## **A Systematic Review of Psychological Distress that Occurs During the Treatment of Tuberculosis (<https://www.psychosocial.com/article/PR270727/18662/>)**

**Authors:** Gevi Melliya Sari, Muhammad Amin, Laily Hidayati

**DOI:** 10.37200/IJPR/V24I7/PR270727 (<https://doi.org/10.37200/IJPR/V24I7/PR2700727>)

**Pages:** 7587-7598

**Keywords:** Anxiety; Depression; Loneliness; Psychological Distress; Tuberculosis

Abstract (<https://www.psychosocial.com/article/PR270727/18662/>)

Download (</register-login>)

## **Factors Associated with Obesity in Adults in South East Asia (<https://www.psychosocial.com/article/PR270728/18663/>)**

**Authors:** Sari M. Ni Ketut Putri, Meo. Christina M, Suhardin. Saverinus, Halimatunnisa M, Inta Susanti

**DOI:** 10.37200/IJPR/V24I7/PR270728 (<https://doi.org/10.37200/IJPR/V24I7/PR2700728>)

**Pages:** 7599-7607

**Keywords:** Obesity, adults, Southeast Asia, risk of obesity

Abstract (<https://www.psychosocial.com/article/PR270728/18663/>)

Download (</register-login>)

## **Effectiveness of Honey Dressing on Healing of Diabetic Foot Ulcer: Systematic Review (<https://www.psychosocial.com/article/PR270729/18664/>)**

**Authors:** Farida Farida, Angga Miftakhul Nizar, Wa Ode Nurlina, Restiyana Agus

**DOI:** 10.37200/IJPR/V24I7/PR270729 (<https://doi.org/10.37200/IJPR/V24I7/PR2700729>)

**Pages:** 7608-7619

**Keywords:** Wound Management; Honey Dressing; Modern Wound Care

Abstract (<https://www.psychosocial.com/article/PR270729/18664/>)

Download (</register-login>)

## **Individual Coaching on Self-Efficacy, Control and Medication Adherence in Patients with Tuberculosis (<https://www.psychosocial.com/article/PR270730/18665/>)**

**Authors:** Tintin Sukartini, Ika Nur Pratiwi, Maria Fatima Koa

**DOI:** 10.37200/IJPR/V24I7/PR270730 (<https://doi.org/10.37200/IJPR/V24I7/PR2700730>)

**Pages:** 7620-7627

**Keywords:** Individual Coaching; Tuberculosis; Self-Efficacy; Adherence; Tuberculosis Treatment

Abstract (<https://www.psychosocial.com/article/PR270730/18665/>)

Download (/register-login)

### **Determinants of the Caregiver Burden of CKD Patients Undergoing Hemodialysis (<https://www.psychosocial.com/article/PR270731/18666/>)**

**Authors:** Virgianti Nur Faridah, Nursalam Nursalam, Ni Luh Putu Inca Buntari Agustini, Trijati Puspita Lestari, Suratmi Suratmi, Farida Juanita, Arifal Aris

**DOI:** 10.37200/IJPR/V24I7/PR270731 (<https://doi.org/10.37200/IJPR/V24I7/PR2700731>)

**Pages:** 7628-7634

**Keywords:** Caregiver; Burden; Hemodialysis; Chronic Kidney Disease

Abstract (<https://www.psychosocial.com/article/PR270731/18666/>)

Download (/register-login)

### **The Relationship Between Spirituality and the Depression Level of the Elderly at the Nursing Home: A Study from Indonesia (<https://www.psychosocial.com/article/PR270732/18677/>)**

**Authors:** Ali Sairozi, Ah. Yusuf, Rizqi Ameliyah, Arival Aris, Iswatun

**DOI:** 10.37200/IJPR/V24I7/PR270732 (<https://doi.org/10.37200/IJPR/V24I7/PR2700732>)

**Pages:** 7635-7640

**Keywords:** Spirituality; Elderly; Depression

Abstract (<https://www.psychosocial.com/article/PR270732/18677/>)

Download (/register-login)

### **Work Stress and Spirituality in Diabetes Mellitus Self-Management (<https://www.psychosocial.com/article/PR270733/18678/>)**

**Authors:** Kusnanto Kusnanto, Nyuasthi Genta Satriyaningarum, Ika Nur Pratiwi, Hidayat Arifin

**DOI:** 10.37200/IJPR/V24I7/PR270733 (<https://doi.org/10.37200/IJPR/V24I7/PR2700733>)

**Pages:** 7641-7647

**Keywords:** work stress; spirituality; diabetes mellitus; self-management

Abstract (<https://www.psychosocial.com/article/PR270733/18678/>)

Download (/register-login)

### **Analysis of Factors Affecting Resilience among Breast Cancer Patients (<https://www.psychosocial.com/article/PR270734/18679/>)**

**Authors:** Lilik Supriati, I Ketut Suidiana, Eka Nurwahyuni, Sri Poeranto, Anis Rosyiatul Husna, Rindayati

**DOI:** 10.37200/IJPR/V24I7/PR270734 (<https://doi.org/10.37200/IJPR/V24I7/PR2700734>)

**Pages:** 7648-7654

**Keywords:** Breast cancer, Resilience

Abstract (<https://www.psychosocial.com/article/PR270734/18679/>)

Download (/register-login)

### **Factors Associated with Headaches in Smartphone Users (<https://www.psychosocial.com/article/PR270735/18680/>)**

**Authors:** Erna Dwi Wahyuni, Aulathivali Inas Faravida, Ika Nur Pratiwi

**DOI:** 10.37200/IJPR/V24I7/PR270735 (<https://doi.org/10.37200/IJPR/V24I7/PR2700735>)

**Pages:** 7655-7662

**Keywords:** duration, frequency, headaches, radiation level, smartphone

Abstract (<https://www.psychosocial.com/article/PR270735/18680/>)

Download (/register-login)

### **How A Team Games Tournament Method Improves Injury Handling Behavior among Those who Participate in Extracurricular Sports (<https://www.psychosocial.com/article/PR270736/18681/>)**

**Authors:** Erna Dwi Wahyuni, Regina Dwi Fridayanti, Lailatun Ni'mah

**DOI:** 10.37200/IJPR/V24I7/PR270736 (<https://doi.org/10.37200/IJPR/V24I7/PR2700736>)

**Pages:** 7663-7670

**Keywords:** Team Games Tournament, Sports Extracurricular Members, Sports Injuries

Abstract (<https://www.psychosocial.com/article/PR270736/18681/>)

Download (/register-login)

### **A Systematic Review of the Impact of Diabetes Resilience on Quality of Life (<https://www.psychosocial.com/article/PR270737/18682/>)**

**Authors:** Hidayat Arifin, Kusnanto Kusnanto, Ika Yuni Widyawati

**DOI:** 10.37200/IJPR/V24I7/PR270737 (<https://doi.org/10.37200/IJPR/V24I7/PR2700737>)

**Pages:** 7671-7678

**Keywords:** Type 2 Diabetes Mellitus; Resilience; Quality of Life

Abstract (<https://www.psychosocial.com/article/PR270737/18682/>)

Download (/register-login)

### **Prevention of Depression in Patients with Cancer: A Systematic Review (<https://www.psychosocial.com/article/PR270738/18683/>)**

**Authors:** Yulia Indah Permata Sari, Tintin Sukartini, Esti Yunitasari

**DOI:** 10.37200/IJPR/V24I7/PR270738 (<https://doi.org/10.37200/IJPR/V24I7/PR2700738>)

**Pages:** 7679-7686

**Keywords:** Anxiety; Antidepressant; Cancer; Depression; Preventive



Abstract (<https://www.psychosocial.com/article/PR270738/18683/>)

Download (/register-login)

**The Correlation between Self-Regulation and Intention with the Diet Adherence of Adults with Type 2 Diabetes Mellitus (<https://www.psychosocial.com/article/PR270739/18684/>)**

**Authors:** Kusananto Kusananto, Ayu Septia Malinda, Khoridatul Bahiyah, Hidayat Arifin

**DOI:** 10.37200/IJPR/V24I7/PR270739 (<https://doi.org/10.37200/IJPR/V24I7/PR2700739>)

**Pages:** 7687=7695

**Keywords:** Adherence; Intention; Self-Regulation; Diabetes Mellitus

Abstract (<https://www.psychosocial.com/article/PR270739/18684/>)

Download (/register-login)

**Determinants of Quality of Nursing Work Life: A Systematic Review (<https://www.psychosocial.com/article/PR270740/18685/>)**

**Authors:** Dluha Maf'ula, Nursalam, Nursalam, Tintin Sukartini

**DOI:** 10.37200/IJPR/V24I7/PR270740 (<https://doi.org/10.37200/IJPR/V24I7/PR2700740>)

**Pages:** 7696-7708

**Keywords:** QNWL; Cultural; Organization Factors; Determinant Factors

Abstract (<https://www.psychosocial.com/article/PR270740/18685/>)

Download (/register-login)

**Undergraduate Nursing Students' Knowledge, Attitude and Practice Toward Palliative Care in Indonesia: A Cross-sectional Online Survey (<https://www.psychosocial.com/article/PR270741/18686/>)**

**Authors:** Ni Luh Putu Inca Buntari Agustini, Nursalam Nursalam, Made Rismawan, Virgianti Nur Faridah

**DOI:** 10.37200/IJPR/V24I7/PR270741 (<https://doi.org/10.37200/IJPR/V24I7/PR2700741>)

**Pages:** 7709-7717

**Keywords:** Knowledge, Attitude, Practice, Undergraduate Nursing Students, Palliative Care

Abstract (<https://www.psychosocial.com/article/PR270741/18686/>)

Download (/register-login)

**A Systematic Review of the Family Acceptance of a Schizophrenic Patient (<https://www.psychosocial.com/article/PR270742/18697/>)**

**Authors:** Darni Darni, Ah Yusuf, Rr Dian Tristiana

**DOI:** 10.37200/IJPR/V24I7/PR270742 (<https://doi.org/10.37200/IJPR/V24I7/PR2700742>)

**Pages:** 7718-7726

**Keywords:** Family; Acceptance; Schizophrenia

**Javanese Family Resilience in Taking Care of Mental Disorder Clients After Pasung**  
(<https://www.psychosocial.com/article/PR270884/19235/>)

**Authors:** Sitti Sulaihah, Ah Yusuf, Hanik Endang Nihayati

**DOI:** 10.37200/IJPR/V24I7/PR270884 (<https://doi.org/10.37200/IJPR/V24I7/PR270884>)

**Pages:** 8971-8978

**Keywords:** Family Resilience; Family's Culture; Javanese; Mental Disorder Clients

Abstract (<https://www.psychosocial.com/article/PR270884/19235/>)

Download (/register-login)

**Resilience in Families of Children with Special Needs from Sibling's Perspective**  
(<https://www.psychosocial.com/article/PR270885/19236/>)

**Authors:** Retno Indarwati, Ika Lusdiana, Eka Mishbahatul, Sylvia Dwi Wahyuni, Rizki Fitryasari, Apriana Rahmawati

**DOI:** 10.37200/IJPR/V24I7/PR270885 (<https://doi.org/10.37200/IJPR/V24I7/PR270885>)

**Pages:** 8979-8986

**Keywords:** Family resilience, sibling, children with special needs

Abstract (<https://www.psychosocial.com/article/PR270885/19236/>)

Download (/register-login)

**Online Games Addiction and the Decline in Sleep Quality of College Student Gamers in the Online Game Communities in Surabaya, Indonesia**  
(<https://www.psychosocial.com/article/PR270886/19237/>)

**Authors:** Andhika Cahya Kharisma, Rizki Fitryasari, Praba Diyan Rahmawati

**DOI:** 10.37200/IJPR/V24I7/PR270886 (<https://doi.org/10.37200/IJPR/V24I7/PR270886>)

**Pages:** 8987-8993

**Keywords:** Online game addiction; Sleep quality; College students; Gamers

Abstract (<https://www.psychosocial.com/article/PR270886/19237/>)

Download (/register-login)

**The Relationship Between Self-Concept, Role of Peers and Family Function and Adolescents' Premarital Sexual Behavior (<https://www.psychosocial.com/article/PR270887/19243/>)**

**Authors:** Praba Diyan Rachmawati, Mira Triharini, Dinda Salmahella

**DOI:** 10.37200/IJPR/V24I7/PR270887 (<https://doi.org/10.37200/IJPR/V24I7/PR2700887>)

**Pages:** 8994-9001

**Keywords:** premarital sexual behavior, self-concept, peer role, family function

Abstract (<https://www.psychosocial.com/article/PR270887/19243/>)

Download (/register-login)

**Prevalence of Mental Disorders in The Community Lamongan-Indonesia: Results of Early Detection with The Community Mental Health Nursing (CMHN) Approach (<https://www.psychosocial.com/article/PR270888/19244/>)**

**Authors:** Arifal Aris, Ah. Yusuf, Virgianti Nur Faridah, Siti Sholihah, Abdul Rokhman, Suhariyati Suhariyati, Ali Sairozi

**DOI:** 10.37200/IJPR/V24I7/PR270888 (<https://doi.org/10.37200/IJPR/V24I7/PR2700888>)

**Pages:** 9002-9007

**Keywords:** Mental disorders; Early detection; CMHN

Abstract (<https://www.psychosocial.com/article/PR270888/19244/>)

Download (/register-login)

**The Role of the Husbands via the Endorphin Massage on the Occurrence of Postpartum Blues in Primiparous Postpartum Mothers (<https://www.psychosocial.com/article/PR270889/19245/>)**

**Authors:** Emuliana Sulpat, Agus Sulistyono, Ah. Yusuf, Andri Tri Kusumaningrum, Susilo Hariyanto, Amellia Mardhika

**DOI:** 10.37200/IJPR/V24I7/PR270889 (<https://doi.org/10.37200/IJPR/V24I7/PR2700889>)

**Pages:** 9008-9013

**Keywords:** husband's role, endorphin massage, postpartum blues.

Abstract (<https://www.psychosocial.com/article/PR270889/19245/>)

Download (/register-login)

**Improving the Cooperative level of Preschool Children During the Intra vena Injection Through Virtual Reality Cartoon Education (<https://www.psychosocial.com/article/PR270890/19246/>)**

**Authors:** Yuni Sufyanti Arief, Regina Soares Da Costa Ximenes, Elida Ulfiana

**DOI:** 10.37200/IJPR/V24I7/PR270890 (<https://doi.org/10.37200/IJPR/V24I7/PR2700890>)

**Pages:** 9014-9019

**Keywords:** Virtual Reality; Cartoons; Education; Cooperative Level, Preschool

Abstract (<https://www.psychosocial.com/article/PR270890/19246/>)

Download (/register-login)

**Risk Factors for Stunting and Severe Stunting among Children (12-59 Months Old) in the Coastal Area of Surabaya, East Java, Indonesia (<https://www.psychosocial.com/article/PR270891/19247/>)**

**Authors:** Eka Mishbahatul Mar'ah Has, Ridha Cahya Prakhasita, Ilya Krisnana, Sylvia Dwi Wahyuni, Yuni Sufyanti Arief, Nursalam Nursalam

**DOI:** 10.37200/IJPR/V24I7/PR270891 (<https://doi.org/10.37200/IJPR/V24I7/PR2700891>)

**Pages:** 9020-9027

**Keywords:** Stunting; Children; Indonesia

Abstract (<https://www.psychosocial.com/article/PR270891/19247/>)

Download (/register-login)

**Social Support and the Implementation of the Early Initiation of Breastfeeding (IMD) among the Madurese in Indonesia (<https://www.psychosocial.com/article/PR270892/19258/>)**

**Authors:** Esti Yunitasari, Pipit Pitaloka, HerdinaMariyanti, Yeo KeeJiar

**DOI:** 10.37200/IJPR/V24I7/PR270892 (<https://doi.org/10.37200/IJPR/V24I7/PR2700892>)

**Pages:** 9028-9034

**Keywords:** Breastfeeding, Early Initiation of Breastfeeding, Social Support

Abstract (<https://www.psychosocial.com/article/PR270892/19258/>)

Download (/register-login)

**Contributing Factors of Personal Protection Equipment (PPE) Utilization among Sand and Gravel (SSG) Mine Workers (<https://www.psychosocial.com/article/PR270893/19259/>)**

**Authors:** Rr Dian Tristiana, Ratih Pravitasari, Erna Dwi Wahyuni

**DOI:** 10.37200/IJPR/V24I7/PR270893 (<https://doi.org/10.37200/IJPR/V24I7/PR2700893>)

**Pages:** 9035-9041

**Keywords:** Attitude; Knowledge; Personal Protection Equipment; Sand and Gravel Mine Workers

Abstract (<https://www.psychosocial.com/article/PR270893/19259/>)

Download (/register-login)

**Nurse's Individual Factors May Predict Quality of Nursing Work Life (Qnwl) in Clinical Setting (<https://www.psychosocial.com/article/PR270894/19260/>)**

**Authors:** Riyanto Faizin, Rizki Fitryasari, Erna Dwi Wahyuni, Nursalam Nursalam

**DOI:** 10.37200/IJPR/V24I7/PR270894 (<https://doi.org/10.37200/IJPR/V24I7/PR2700894>)

**Pages:** 9042-9049

# Determinants of the Caregiver Burden of CKD Patients Undergoing Hemodialysis

Virgianti Nur Faridah<sup>1,2</sup>, Nursalam Nursalam<sup>1</sup>, Ni Luh Putu Inca Buntari Agustini<sup>1</sup>, Trijati Puspita Lestari<sup>2</sup>, Suratmi Suratmi<sup>2</sup>, Farida Juanita<sup>2</sup>, Arifal Aris<sup>2</sup>

**Abstract** -- Families who treat chronic kidney disease (CKD) patients are at risk of experiencing caregiver burden. The family caregiver burden depends on the ability of the families to care for their family members who experience CKD when undergoing hemodialysis. The purpose of this study was to determine the factors associated with the caregiver burden of caring families (primary caregivers). This study used quantitative research and a cross-sectional design. The sample consisted of 95 respondents determined using consecutive sampling from the population in the Hemodialysis room of Muhammadiyah Lamongan Hospital. The variables of this study were caregiver burden (Y), Age (X1), Gender (X2), Educational level (X3), Salary (X4) and Treating the patient's time (X4). The research instrument consisted of the Zarit Burden Interview (ZBI) method to determine the caregiver burden. Multivariate analysis was done using logistic regression. The results showed that the caregiver model shows a significant burden with a variable in salary and age. The accuracy of the model when predicting caregiver burden was 62.1%. These two variables can explain the burden of the caregivers by 10%. The equation model obtained the results of Caregiver burden =  $-0,277 + 0,818 * \text{salary} - 1,313 * \text{age (adults)} - 0,738 * \text{age (elderly)}$ . There is a need for social support from other families who care. Support system can influence the level of caregiver stress.

**Keywords** – Caregiver; Burden; Hemodialysis; Chronic Kidney Disease

---

## I. INTRODUCTION

Chronic kidney disease and hemodialysis therapy is a burdensome and complex therapy that requires a lot of support from the family [1]. Hemodialysis can reduce the patient's energy level and affect the patient's ability to work and perform daily activities. This disrupts the normal life of the patient and their family [2]. The families often experience a tremendous burden associated with caring for hemodialysis patients.

Jafari et al research showed that 42.7% of families had a moderate level of care burden, 32.5% had a high level, and 4.9% had a very high level [3]. A systematic review by Gilbertson et al concluded that the burden and quality of life of families where a family member was on hemodialysis was worse than the general population. This is comparable to other chronic diseases. The burden of care for hemodialysis patients is comparable to dialysis peritoneal patients. The quality of life of families post-kidney transplant is better than that of dialysis patients [1]. The results of the 2 studies are

---

<sup>1</sup>Faculty of Nursing, Universitas Airlangga Surabaya, Indonesia

<sup>2</sup>Faculty of Health Science, Universitas Muhammadiyah Lamongan, Indonesia

Corresponding author  
Trijati Puspita Lestari  
Email: [pipid.puspita@gmail.com](mailto:pipid.puspita@gmail.com)

different because they are viewed from different points of view. It can still be concluded that families experience a significant burden when treating patients with hemodialysis.

The caregiver or family challenges and burdens increase with the progression of the disease. Families experience physical burdens were associated with a lack of social support and lower health-related quality of life for both the patients and their families [5]. Disorders related to quality of life and family burden in the hemodialysis patients places a double burden on them which interferes with the treatment process [3].

Several factors affect the burden of care such as the level of the patient's ability to care for themselves (self-care), chronic comorbidities, level of education and the patient's family age. If the patient's self-care ability is low, there is an incident of chronic disease accompanying the patient, a low level of education and increasing family age, then the level of care burden on the family will increase [3]. Care burden is a term used by the caregivers or their families to refer to the hardships experienced when caring for the patients [6]. This burden covers the physical, psychological, social, and financial aspects. Increased care burden and decreased quality of life can cause complications such as depression. There is also a significant relationship between increasing the burden of care and decreasing the ability of care provided by the family because the burden of care can greatly impact an individual. The amount of stress experienced by the family due to the experience of caring for patients with a chronic illness is a serious, hidden disease [6]. The burden of care affects the quality of life of the family and it results in the provision of care being reduced. Thus, the condition of patients with a chronic illness worsens. The worsening of the patient's condition can increase the burden of care and cause a repetitive cycle. If a timely intervention is not carried out, then it can cause gradual fatigue in the family [3].

The purpose of this study was to determine the factors associated with caregiver burden regarding patients with hemodialysis. We can use this to determine the direction of advanced care planning. Family-centered care has been proposed to address the problem by meeting the needs of not only the patients but also their family members [7]. Efforts undertaken to prepare the families for the care of hemodialysis patients include more family-centered care, the provision of appropriate education early on, and interventions that target the readiness of the care partners. This is in addition to communication that focuses on the state of readiness between the health care providers and the patients' family members, and the establishment of a dynamic relationship between the family and patient [8]. The formation of psycho-educational support groups can also empower the families by providing the knowledge and emotional support needed for successful care [9].

## **II. METHODS**

This study used quantitative research and a cross-sectional design [10]. The variables of this study were caregiver burden (Y), age (X1), gender (X2), educational level (X3), salary (X4) and treating the patient's time (X4). The research instruments consisted of the Zarit Burden Interview (ZBI) to determine the caregiver burden [11], [12]. Multivariate analysis was done using logistic regression.

- Data collection

The study subjects were taken on a consecutive basis from those who meet the criteria. The sample size was calculated using sample size software. From the calculation results, we obtained a minimum sample size of 95 respondents from the Hemodialysis room of Muhammadiyah Lamongan Hospital.

- Sample and Sampling

The sample consisted of the primary caregivers of CKD patients who had undergone treatment for hemodialysis. The criteria for the caregivers is that they have directly cared for CKD patients for at least one month after the patient has undergone hemodialysis while not caring for other patients for a long and concurrent time.

- Instruments

The research instrument was the Zarit Burden Interview (ZBI) to determine the caregiver burden. The Zarit Burden Interview (ZBI) method was used to explore the extent of the caregiver burden. It contains 22 items used for investigating the 5 domains of caregiver burden. The questionnaire items were rated using a 5-point Likert scale with 0 (rarely) being the lowest and 4 (nearly al-ways) being the highest. The total score can explain the level of caregiver burden. A score of 0–20 indicates little or no burden, 21–40 is a mild to moderate burden, 41–60 is moderate to severe burden and 61–88 is a severe burden is present [11], [12].

### III. RESULTS

- General Characteristics of the Participants

The demographic data of the respondents (Table 1) showed that the caregivers of the patients with CKD undergoing hemodialysis were made up of 48,4% elderly (more than 45 years old), 60% female; 64,2% graduates of senior high school; 56,8% with a salary > 2 million/month and 43,2% were treating the patient for more than > 3 years. The studies showed that the caregivers of the patients with hemodialysis were mostly women and elderly people [8], [12], [13], [14], [15]. The studies conducted by Jafari, et al showed that the caregivers of patients with CKD had a high school level of education [3], [15]. Income is categorized based on the Lamongan Regency minimum wage, which is low if it is less than 2.2 million, and high if it is more than 2.2 million per month [16].

Table 1. General characteristics (N=95)

Demographic data	Σ	%	Mean	SD
Age			2.33	7.36
Teenagers (< 25 <sup>th</sup> )	15	15.8		
Adult (25 <sup>th</sup> – 45 <sup>th</sup> )	34	35.8		
Elderly (> 45 <sup>th</sup> )	46	48.4		
Total	95	100		
Gender			1.40	0.496
Female	57	60		
Male	38	40		
Total	95	100		
Education level			2.98	0.601
Elementary school	0	0		
Junior high school	18	18.9		
Senior high school	61	64.2		
University	16	16.8		
Total	95	100		
Salary			1.57	0.498
< 2 million/month	41	43.2		
> 2 million/ month	54	56.8		
Total	95	100		
Treating patient's time			2.22	0.774
< 1th	20	21.1		
1-3th	34	35.8		
>3th	41	43.2		
Total	95	100		

- Descriptive Analysis of ZBI

The family burden when treating patients with CKD undergoing hemodialysis is in the category of no burden or low burden for 55 families (57.9%). There burden was felt by 40 families (42.1%). This does not follow Jafari's research showing that 42.7% of families had a moderate level of care burden, 32.5% had a high level, and 4.9% had a very high level [3]. The differences in these findings can be attributed to the culture of Indonesians who care for their family members sincerely. This is in addition to the differences in the research environment and the number of supporting facilities available for the caregiver to access. However, the family care burden is still high at 42,1%. This indicates the need for attention and solutions.

- Chi square test

The variables assessed using the Chi-square test were age, sex, education, economy (salary), and the length of care of the patients. The variable that will be removed is any variable with a  $p > 0.250$ . The Chi-square test results showed that the sex variable had a  $p = 0.525$ , so that variable was not included in the logistic regression test.

Table 2. Chi square test results

Demographic data	Chi square
Age	0.134
Gender	0.525
Educational level	0.057
Salary	0.117
Treating patient's time	0.127

- Logistic Regression Test

The variables tested using logistic regression were age, educational level, salary, and treating the patient's time. Treating the patient's time and the education level variables were not significant in relation to the caregiver burden variable with  $p = 0.034$  and  $p = 0.023$ . These variables were removed from the model. Based on this research, salary is the determinant factor of caregiver burden.

The caregiver burden model is significant in relation to the variables of salary and age. The accuracy of the model when predicting caregiver burden is 62.1%. These 2 variables can explain the caregiver burden by 10%. The model equation obtained is:

$$\text{Caregiver burden} = -0,277 + 0,818 * \text{salary} - 1,313 * \text{age (adults)} - 0,738 * \text{age (elderly)}$$

Based on the caregiver burden model, families with an economy of <2 million / month have a risk of experiencing a burden when treating patients as much as 2,262 times compared to families with an economy of > 2 million / month. Families in adulthood (25-45 years old) are at risk of experiencing a burden when caring for patients of 0.269 compared to the families of adolescents (<25 years). Elderly families (> 45 years old) are at risk of experiencing a burden when caring for patients of 0.478 compared to the families of adolescents (Table 3).

Table 3. Caregiver Burden Model

	B	S.E.	Wald	Df	Sig.	Exp(B)
Step 1 <sup>a</sup>						
Salary(1)	.816	.445	3.364	1	.067	2.262
Age			4.781	2	.092	
Age(1)	-1.313	.679	3.735	1	.053	.269
Age(2)	-.738	.475	2.411	1	.120	.478
Constant	-.227	.344	.434	1	.510	.797

a. Variable(s) entered in Step 1: Salary, Age.



#### **IV. DISCUSSION**

Family members are often involved in the treatment of chronic kidney disease patients as their care partners. As care partners, the family members are involved in many important activities, some of which include administering medication, overseeing medication compliance, assisting in daily care activities, scheduling and providing transportation when attending medical appointments, monitoring patient health, advocating for the patients and offering support emotionally [5]. The role of the family is beneficial for both the patients and health care providers. For patients, the involvement of their family members is associated with improved behavior, self-management care and adherence to therapy, in addition to improving their quality of life, reducing the risk of death, reducing anxiety and depressive symptoms, and decreasing the hospital admission opportunities. For the healthcare providers, the family members are invaluable partners in the task of caring for patients with chronic kidney disease [8].

The caregivers of patients on hemodialysis are at a high risk of emotional development, psychological distress, low quality of life, and increased burden. They have a variety of psychosocial concerns because of the length of the treatment. They usually experience an increased workload, limited personal and social activities, and financial problems. They feel tired, angry, depressed, helpless, guilty, isolated, that they are losing their freedom, fearful, vulnerable, and that they are neglecting their health. Interventions and supportive information are very important for the caregivers because they can help them to improve their quality of life and their ability to cope effectively with the care of their patients. Caregivers need appropriate knowledge, special skills, education, and guidance to help their patients to achieve better adherence to hemodialysis treatment [17].

The family caregivers of palliative care sufferers often experience enormous physical burdens in addition to the emotional and financial responsibilities associated with providing care. The caregiver burden increases over time non-linearly from the initial diagnosis to the death of the patient [18]. Care for the family caregivers (Family Care Giver) is needed in all domains of quality of life (physical, psychological, social and spiritual) [9].

The caregiver burden model is significant in relation to the variables of salary and age. This is relevant to the study stating that caregivers with an inappropriate economic status are significantly more prone to care burden [2]. Mashayekhi et al. found there to be a statistically significant relationship between their income and the level of care burden felt by the caregiver. Thus the caregivers of lower-income patients experience a higher level of care load [19]. The conceptual model of caregiver burden expressed by Gabriel is based on Lazarus and Folkman's theories, which show that an individual's response to stress is influenced by the primary appraisal, secondary appraisal, and available resources. According to this model, primary appraisal, secondary appraisal, and the availability of interacting resources ultimately have an impact on the individual stress response. Caregivers assess the effects of various aspects of care on themselves in the primary appraisal, which includes potential factors for patients and caregivers. The caregiver characteristics include caregiver demographics, employment status and family income. Transactions between the external resources in the form of social support and internal resources in the form of self-efficacy can affect the burden of care [20].

Bayoumi reported there to be a positive and significant correlation between total care burden and patient age. There was also a significant but negative correlation between total care burden and nurse age, patient education level and

caregiver education level. The difference in our findings and the results of the study above could be due to the low sample size (50 caregivers) in the Bayoumi study because the low sample size reduces the level of study transferability and the opportunity to find any correlations and relationships between the variables [21]. Several factors affect the burden of care such as the level of the patient's ability to care for themselves (self-care), chronic comorbidities, level of education and the patient's family age. If the patient's self-care ability is low, there is an incident of chronic disease accompanying the patient, a low level of education and an increased family age, then the level of care burden felt by the family will increase [3].

A comprehensive understanding of the family carers' unmet needs is needed to develop a long-term care plan for the family caregivers [22]. The palliative system must take on a more active role involving patients and their families as partners in order to manage the treatments and side effects and to achieve the best patient outcomes. They must be able to evaluate the patients and caregivers in order to determine what tasks they will be able to do before making sure that they have the training and resources to carry out these tasks [23].

## V. CONCLUSION

The caregiver model is a significant burden with a variable in relation to salary and age. The determinant factor of caregiver burden for CKD patients is the economic factors. Families with a poor economic status will be more vulnerable to the burden of care. The caregivers of low-income patients experience higher levels of care load. A person's burden can be removed by the utilization of available resources. The role of the family members is needed to reduce the caregiver's burden.

## CONFLICT OF INTEREST

No conflicts of interest have been declared.

## ACKNOWLEDGMENT

The authors would like to acknowledge the Faculty of Nursing in Universitas Airlangga, Surabaya, especially the nursing doctoral programs for the support granted to the researchers. They would also like to thank the Director of Muhammadiyah Lamongan Hospital and Rector Universitas Muhammadiyah Lamongan.

## REFERENCES

- [1] E. L. Gilbertson, R. Krishnasamy, C. Foote, A. L. Kennard, M. J. Jardine, and N. A. Gray, "Burden of Care and Quality of Life Among Caregivers for Adults Receiving Maintenance Dialysis: A Systematic Review," *American Journal of Kidney Diseases.*, vol. 73, no. 3, pp. 332–343, 2019.
- [2] Abbasi A, Rahmani H, Shariati A, Asayesh H, Ashrafrezaee N, Mollaei E, "The relationship between caring burden and coping strategies in hemodialysis patients caregivers," *Journal Urmia Nurs Midwifery Fac.*, vol. 10, no. 4, pp. 533–9, 2012.
- [3] H. Jafari, A. Ebrahimi, A. Aghaei, and A. Khatony, "The relationship between care burden and quality of life in caregivers of hemodialysis patients," *BMC Nephrology*, vol. 19, no. 1, p. 321, 2018.
- [4] V. Maddalena, F. O'Shea, and B. Barrett, "An exploration of palliative care needs of people with end-stage renal disease on dialysis: Family caregiver's perspectives," *Journal Palliative Care*, vol. 33, no. 1, pp. 19–25, 2018.
- [5] V. L. Hoang, T. Green, and A. Bonner, "Informal caregivers' experiences of caring for people receiving dialysis: A mixed-methods systematic review," *Journal of Renal Care*, vol. 44, no. 2, pp. 82–95, 2018.
- [6] E. A. Sajadi SA, "Quality of life among family caregivers of patients on hemodialysis and its relevant factors: a systematic review," *Int J Community Based Nurs Midwifery*, vol. 5, no. 3, p. 206, 2017.

- [7] K. M. Kokorelias, M. A. M. Gignac, G. Naglie, and J. I. Cameron, "Towards a universal model of family centered care : a scoping review," *BMC Health Services Research*, vol. 5, pp. 1–11, 2019.
- [8] N. DePasquale, A. Cabacungan, P. L. Ephraim, L. Lewis-Boyer, N. R. Powe, and L. E. Boulware, "Family Members' Experiences With Dialysis and Kidney Transplantation," *Kidney Medicine*, vol. 1, no. 4, pp. 171–179, 2019.
- [9] et all Ferrell, Betty R, Kravitz, Kate, "Family Caregivers," *Clinical Journal Oncology Nursing.*, vol. 22, no. 3, pp. 286–294, 2018.
- [10] Nursalam, "Metodologi Penelitian Ilmu Keperawatan Pendekatan Praktis", 4th ed. *Jakarta: Salemba Medika*, 2015.
- [11] S. Zarit, K. Reever, and J. Bahc-Peterson, "How Emotions Drive Customer Loyalty (Infographic) | Provide Support," *Gerontologist*, vol. 20, no. 6, pp. 649–655, 1980.
- [12] I. Cantekin, M. Kavurmaci, and M. Tan, "An analysis of caregiver burden of patients with hemodialysis and peritoneal dialysis," *Hemodialysis International.*, vol. 20, no. 1, pp. 94–97, 2016.
- [13] M. Sellars *et al.*, "An Interview Study of Patient and Caregiver Perspectives on Advance Care Planning in ESRD," *American Journal of Kidney Diseases*, vol. 71, no. 2, pp. 216–224, 2018.
- [14] K. Griva *et al.*, "Quality of life and emotional distress in patients and burden in caregivers: a comparison between assisted peritoneal dialysis and self-care peritoneal dialysis," *Quality of Life Research*, vol. 25, no. 2, pp. 373–384, 2016.
- [15] L. Rabiei, A. A. Eslami, H. A. Abedi, R. Masoudi, and G. R. Sharifirad, "Caring in an atmosphere of uncertainty: perspectives and experiences of caregivers of peoples undergoing haemodialysis in Iran," *Scandinavian Journal of Caring Sciences*, vol. 30, no. 3, pp. 594–601, 2016.
- [16] "SK UMK Gubernur Jatim." 2019.
- [17] G. Eirini and G. Georgia, "Caregivers of Patients on Haemodialysis," *Caregiving Home Care*, 2018.
- [18] D. Guerriere, A. Husain, B. Zagorski, D. Marshall, and H. Seow, "Predictors of caregiver burden across the home-based palliative care trajectory in Ontario , Canada," *Health and Social Care in the community*, vol. 24, pp. 428–438, 2016.
- [19] F. Mashayekhi, M. Pilevarzadeh, and F. Rafati, "The Assessment of Caregiver Burden in Caregivers of Hemodialysis Patients," *Materia Socio Medica*, vol. 27, no. 5, p. 333, 2015.
- [20] I. O. Gabriel *et al.*, "Caregiver Burden Among Informal Caregivers of Women with Breast Cancer," *Biomedical Journal of Scientific & Technical Research*, pp. 11384–11392, 2019.
- [21] M. M. Bayoumi, "Subjective Burden on Family Carers of Hemodialysis Patients," *Open Journal Nephrology*, vol. 04, no. 02, pp. 79–85, 2014.
- [22] Y. Kim, C. S. Carver, and A. Ting, "Seminars in Oncology Nursing Family Caregivers ' Unmet Needs in Long-term Cancer Survivorship," *Seminar Oncology Nursing*, vol. 35, no. 4, pp. 380–383, 2019.
- [23] C. W. Given, "Seminars in Oncology Nursing Family Caregiving for Cancer Patients : the State of the Literature and a Direction for Research to Link the Informal and Formal Care Systems to Improve Quality and Outcomes," *Semininar Oncology Nursing*, vol. 35, no. 4, pp. 389–394, 2019.