

RESEARCH ARTICLE

Examining the effects of Lemongrass (*Cymbopogon nardus*) extract on Sperm Quality and Testosterone Level in Mice Induced with *Staphylococcus aureus*

Lilis Maghfuroh¹, Putri Ayuika Setiyowati^{2*}, Rofiatun Solekha²,
Alif'iyah Mar' Atus Sholikhah¹, Angella Ananda Saputra², Eka Febrianti Wulandari¹

¹Nursing Program, Faculty of Health Science, Universitas Muhammadiyah Lamongan, East Java, Indonesia.

²Biology Program, Faculty of Science, Technology, and Education,
Universitas Muhammadiyah Lamongan, East Java, Indonesia.

*Corresponding Author E-mail: putriayuikasetiyowati@gmail.com

ABSTRACT:

There are a variety of causes for male infertility. Among them is microbial infection. Lemongrass (*C. nardus*) contains natural antioxidants and flavonoids that have the ability to preserve the male reproductive system. This study examined the effects of different *C. nardus* extract dosages on mouse sperm quality. The male Balb/C mice were separated into five groups of equal size. The negative control group was injected with 0.01ml of Na-CMC 0.5% over the course of 28 days, the positive control group was injected with 0.2ml of *Staphylococcus aureus* (*S. aureus*) on days 10, 17, and 24, and the treatment groups were injected with 0.2ml of *S. aureus* on days 10, 17, and 24, followed by 0.01ml of *C. nardus*. After the experiment, mice were slaughtered, and sperm suspension was obtained from the cauda epididymal to measure morphology, concentration, and motility. The testis was fixed in a fixative solution to analyze the spermatogenic cells, and intracardiac blood was drawn to determine the level of testosterone. Significant differences (p 0.05) were seen in the morphology, concentration, and motility of spermatozoa when *C. nardus* extract was administered. Given the low and middle concentrations of *C. nardus* extract, it can repair sperm damage caused by *S. aureus* infection.

KEYWORDS: Male infertility, *C. nardus* extract, *S. aureus*, Sperm quality, Testosterone level.

INTRODUCTION:

Infertility is becoming more common in Indonesia every year. Infertility affects not only women but also men. Men are responsible for 36% of fertility success^{1,2}. Male infertility can be caused by a variety of factors, including bacteria pathogens such as *Chlamydia sp.*, *Escherichia coli*, *Mycoplasma hominis*, and *Staphylococcus aureus*. This bacteria has been shown to reduce sperm motility^{3,4,5}. *Staphylococcus aureus* (*S. aureus*) can cause infections in a variety of ecological niches within the host⁶. It colonizes the nares, axillae, vagina, pharynx, and damaged skin surfaces in humans, causing a variety of suppurative (pus-forming) infections and toxinoses^{7,8}.

Many studies have revealed that *S. aureus* can cause male infertility^{9,10}. One study revealed that incubating *S. aureus* in sperm reduced testis weight, seminiferous tubulous diameter, epididymal sperm count, and serum testosterone level^{11,12}. Another study found that *S. aureus* alters the composition of sperm proteins and promotes early cell death by secreting lipopolysaccharides (LPS). It resulted in an increase in dead cells and a decrease in motile sperm cells^{3,13}. In this case, the use of antioxidant-containing plants is thought to reduce the toxic effect of various toxicants, including bacterial infection^{14,15}. The concentration of spermatozoa in one ejaculation depends on the process of formation of spermatogenesis in seminiferous tubules¹⁶. When spermatogenesis proceeds normally it will be generated a normal spermatozoa count, otherwise during the process spermatogenesis occurs interference, then the development of spermatogonia cells will affect the number of spermatozoa formed. It

depends on the magnitude of the disturbance that occurred during the process of spermatogenesis^{16,17}. Lemongrass is a natural antioxidant with numerous benefits that are commonly used in herbal medicine¹⁸. *Cymbopogon nardus*, one type of lemongrass studied in this study, is known for its antioxidant activity and effectiveness to attack bacteria by lysis the cells via its potent flavonoid compound^{19,20}. As a result, this study looked into the potential protective effects of *C. nardus* extract on the male reproductive system after infection with *S. aureus*. The parameters measured were epididymal and testicular weight, the morphology of sperm, concentration of sperm, motility of sperm, spermatogenic cells, the diameter of seminiferous tubules, and testosterone level.

MATERIALS AND METHODS:

Ethical approval:

With the number 120-KEP-UB-2021, the Ethical Review Committee of the Research and Community Service Department at Universitas Brawijaya, East Java, Indonesia, approved all procedures in this study, including the use of mice as animal models.

Plant-based substances:

C. nardus was collected in September 2021 from Materia Medica, Batu, East Java. It was acknowledged and authenticated by Universitas Muhammadiyah Lamongan's Biological Laboratory, Biology Department.

Na-CMC 0.5% preparation and suspension:

A total of 500 milligrams of *Na-Carboxymethyl cellulose* (Na-CMC) were weighed and dissolved in warm distilled water. Using a 100ml measuring flask, all of the remaining distilled water was added after dissolving to obtain a volume of 100ml Na-CMC solution.

C. nardus ethanolic extract preparation and suspension:

C. nardus stalks were cut into little pieces and dried in the sun for about five days. The 250g of dried stalks were ground in an electric grinder before being macerated with ethanol at room temperature for three days. Then, the extract was filtered and concentrated in a water bath at 70⁰ degrees Celsius. To create the suspension, the extract was homogeneously mashed with 0.5% Na-CMC. The respective concentrations were 0.5%, 1%, and 2%.

Microorganism:

The strain of *S. aureus* employed in this investigation was previously obtained from the cervix of an infertile woman in our laboratory.

Substances and chemicals:

Ethanol, methanol, eosin, nigrosine, chloroform (purchased from Merck, Darmstadt, Germany), distilled water, formalin 10% neutral buffer, paraffin, hematoxyline eosin, testosterone kit, xylol, alcohol 70%, alcohol 96%, and NaCl 0.9%.

Animals:

Thirty mature male Balb/C mice were donated from the Veterinary and Pharma Center in Surabaya, East Java (eight weeks old, weighing 25–30grams). They were fed and watered ad libitum in standard laboratory conditions (temperature room at 28–30°C, and 12-h–12-h light-dark cycle).

Experimental design:

After a week of acclimatization, the animals were put into five equal groups (n = 6). The negative control groups were given 0.1ml of Na-CMC 0.5% subcutaneously every day for 28 days. The positive control groups were given 0.2ml of *S. aureus* (0.5 Mc Farland) three times on days 10, 17, and 24. As positive controls, 0.2ml *S. aureus* (0.5McFarland) was given three times to the treatment groups through the intraperitoneal route. Then, different doses of 0.1ml were given subcutaneously to each treatment group in a steady stream. The first group was given 25mg/kg of body weight, the second group was given 50mg/kg of body weight, and the third group was given 100mg/kg of body weight. All three treatments were given over the duration of 28 days. At the end of the process, chloroform was used to kill all of the mice. Then, the cauda epididymal was taken and used to make a sperm suspension for measuring sperm.

Sperm analysis:

This study initially measured epididymal and testicular weight from each sample, then measured the percentage of motility, morphology, and the concentration of spermatozoa in a suspension. 10µL was evaluated under a light microscope in a hemocytometer chamber for sperm motility. One hundred sperm cells were counted in each animal, and the percentage of motility was classified as motil (%) and immotile (%). The motile sperms have moved straight. An hour after sperm diffusion in the solution, 10µL of the sperm suspension was transferred to each hemocytometer counting chamber for 5 minutes to evaluate sperm concentration. Using a 40-magnification light microscope, sperm heads were counted and quantified as million/ml of suspension. Eosin-nigrosine staining determined sperm morphology. To make smears, spermatozoa suspension is dropped on the object's glass and dried in the air. Smears were fixed with methanol, colored with 1% Y eosin solution and nigrosine, and dried. Distilled water cleaned and dried the preparations. 100 mouse

spermatozoa were examined using a 400-magnification light microscope. Calculating the normal/abnormal spermatozoa percentage was the final stage.

Levels of Testosterone:

Serum was taken from intracardiac blood. An enzyme-linked immunoassay kit (Bioassay Technology Lab® 96 kit testosterone, China) evaluated serum testosterone. The level of testosterone measured as nmol/mL

Examining Spermatogenic Cells Histologically:

After fixing the testicular collection in neutral buffer formalin 10%, paraffin, and hematoxyline eosin were used to make histological preparations with a 5µm thickness. The diameter of seminiferous tubules and the number of spermatogenic cells (spermatogonia, spermatid, and spermatocytes) were examined in mouse testicular preparations under 400x magnification.

Data analysis:

All data were represented as means standard error of the mean (SEM), and statistical analysis was conducted using one-way analysis of variance (one-way ANOVA) and post hoc multiple comparison test to determine differences between pairs of means.

RESULT:

The results showed that giving *S. aureus* to mice for 28 days led to a significant ($p < 0.05$) decrease in the quantity of sperm and the percentage of spermatozoa that could move (immotile). In mice that were infected by *S. aureus*, the percentage of spermatozoa that had an aberrant morphology was statistically ($p < 0.05$) higher than it was in the group that served as a negative control (K1). When compared to animals infected with *S. aureus*, testosterone levels and sperm parameters such as motility and concentration increased significantly ($p < 0.05$) in mice that were given *C. nardus* extract (Table 1). The diameter of the seminiferous tubule, the quantity of spermatogonia cells, spermatocytes, and spermatid

cells in the positive control that was infected by *S. aureus* all had lower values than the normal negative control and the group that was treated with *C. nardus* extract. This was another finding (Table 2).

According to the data in table 1, spermatozoa motility at a dose of 100 mg/kg body weight of *C. nardus* extract (P3) was not significantly different from the positive control group (K2). This means that at high doses, the ability to restore the quality of spermatozoa damaged by lead acetate exposure was not optimal. Furthermore, at a dose of 50 mg/kg body weight, the *C. nardus* extract demonstrated the highest average in each observational parameter. It is reasonable to assume that the dose was optimal for improving spermatozoa quality. A description of sperm morphological observation was shown in Figure 1. The differences in the histological seminiferous tubule in each group were shown in figure 2.




Fig.1: Sperm cell morphology in adult mice had normal (N) and abnormal (A: twisted midpiece or tail; B: head malformation (like a needle); C: larger head; D: broken tail).

Table 1: The effect of *C. nardus* extract on sperm motility, morphology, concentration, and testosterone levels in *S. aureus*-infected mice.

Parameters	K1	K2	P1	P2	P3
Motility (%)	80,60 ± 2,51 ^a	58,20 ± 2,28 ^b	70,40 ± 2,70 ^{ab}	76,20 ± 4,02 ^a	61,00 ± 1,58 ^b
Normal morphology (%)	83,60 ± 2,30 ^a	41,60 ± 2,07 ^{bc}	67,20 ± 1,92 ^{ab}	70,40 ± 3,36 ^{ab}	51,00 ± 2,92 ^b
Concentration (x10 ⁶ / mL)	83,00 ± 2,55 ^a	50,00 ± 2,24 ^{ab}	72,20 ± 1,48 ^b	77,20 ± 1,92 ^{bc}	65,80 ± 2,78 ^c
Testosteron level (nmol/L)	13.68 ± 0.65 ^a	8.60 ± 0.40	10.90 ± 0.57	12.89 ± 0.85 ^a	9.69 ± 0.42
Testicular weight (mg)	306 ± 20,73 ^a	216 ± 20,73 ^{ab}	258 ± 23,76 ^b	292 ± 23,88 ^a	245 ± 7,07 ^{ab,b}
Epididymal weight (mg)	10,84 ± 0,39 ^a	7,68 ± 0,40	10,02 ± 0,58 ^{ab}	10,36 ± 0,63 ^{a,ab}	8,46 ± 0,34

Data are presented as mean ± SD. Differences in superscripts in the same rows show significant differences ($p < 0,05$).

K1 (negative control): Na-CMC 0,5%; K2 (positive control): *S. aureus* 0,5 Mc Farland; P1 (treatment group 1): *S. aureus* 0,5 Mc Farland + *C. nardus* 25 mg/kg BW; P2 (treatment group 2): *S. aureus* 0,5 Mc Farland + *C. nardus* 50 mg/kg BW; P3 (treatment group 3): *S. aureus* 0,5 Mc Farland + *C. nardus* 100 mg/kg BW.




Fig. 2: Testicular seminiferous tubules in different groups. Red arrows indicate spermatogonia cells, yellow spermatocytes, blue spermatids, and black seminiferous tubule diameter. (Hematoxylin Eosin; 400x).

Table 2. Effect of *C. nardus* extract on spermatogenic cells and diameter of the seminiferous tubule in mice infected by *S. aureus*

Parameters	K1	K2	P1	P2	P3
DST	216,52 ± 16,65 ^a	105,60 ± 13,91 ^b	141,41 ± 13,19	186,03 ± 29,70 ^a	115,79 ± 13,85 ^b
Spermatogonia cells	82,80 ± 4,38 ^a	60,00 ± 4,00	75,20 ± 4,60 ^{ab}	80,00 ± 3,16 ^{a,ab}	67,20 ± 4,15
Spermatocytes cells	99,60 ± 5,73 ^a	65,20 ± 7,95	87,80 ± 6,72 ^{ab}	98,00 ± 6,33 ^a	81,20 ± 3,35 ^{ab}
Spermatid cells	284,0 ± 42,9 ^a	179,2 ± 18,6	255,2 ± 39,5 ^{a,ab}	268,0 ± 24,5 ^{a,ab}	204,8 ± 15,1

Data are presented as mean ± SD

Differences in superscripts in the same rows show significant differences ($p < 0.05$). K1 (negative control): Na-CMC 0,5%; K2 (positive control): *S. aureus* 0,5 Mc Farland; P1 (treatment group 1): *S. aureus* 0,5 Mc Farland + *C. nardus* 25 mg/kg BW; P2 (treatment group 2): *S. aureus* 0,5 Mc Farland + *C. nardus* 50 mg/kg BW; P3 (treatment group 3): *S. aureus* 0,5 Mc Farland + *C. nardus* 100 mg/kg BW. DST= Diameter of Seminiferous Tubules

DISCUSSION:

In ethnomedicines, the use of medicinal plants, commonly known as medicinal herbs, has been suggested for the treatment of a wide variety of disorders at various points in history and continued to be practiced today. Throughout the history of humanity, people have relied on the therapeutic properties of various plants to heal a variety of illnesses^{21,22}. New scientific approaches have been integrated into conventional medical practice to facilitate the standardization of phytotherapy, which refers to the use of numerous plant-based remedies in combination with one another^{23,14}.

According to this study, The decreased quality of spermatozoa that has been caused by bacterial infection was partially restored by the use of an extract of *C. nardus*. There was a statistically significant difference

($p < 0.05$) between the group that was treated with *C. nardus* extract and the group that served as the positive control and was treated with *S. aureus*. To acquire the optimal results in terms of epididymal and testicular weight, sperm motility, normal morphology of sperm, testosterone level, sperm concentration, and the number of spermatogenic cells, a dose of 25mg/kg of body weight was shown to be the most effective. The increased dose that was administered did not result in an improvement in the quality of the spermatozoa. Even at a dose of 100mg/kg of body weight, the quality of sperm that had been infected with *S. aureus* was not improved.

S. aureus is a pathogen bacteria that are typically found in the female genital tract². This bacteria is known to have a toxic effect on the epididymal duct in addition to disrupting the process of spermatogenesis and also could well interact with the CD14 on the surface of the sperm, primarily in the head and tail region²⁴. This interaction will result in increased levels of sperm membrane lipid peroxidation, production of reactive oxygen species, and caspase-mediated apoptosis in the subsequent stages^{6,10,25}. A reduction in the motility of spermatozoa and an increase in lipid peroxidation are both indicators that cell damage has been caused by *S. aureus*^{26,16,27}

The process of male reproduction is a complicated one that requires a great deal of energy. The hypothalamus and the pituitary gland use a variety of feedback regulation mechanisms to control the function of the testis, which serves as the primary organ of the male germline^{1,28}. GnRH, FSH, and LH are the primary hormones responsible for controlling regulatory mechanisms via their actions on the hypothalamus-pituitary-testis axis²⁹. Through neurons that secrete GnRH, the hypothalamus can monitor the condition of the reproductive system. Through the anterior pituitary, GnRH stimulates the release of LH as well as FSH. LH stimulates the production of testosterone by acting on testicular interstitial cells and increasing their permeability. FSH is the hormone that is responsible for the stimulation of the seminiferous tubules, which is necessary to maintain sperm production³⁰.

A decrease in the amount of testosterone in the body can lead to reproductive disorders. Testosterone is a hormone that is primarily found in men³¹. The precise and coordinated secretion of GnRH, LH, FSH, and testosterone is what regulates male reproductive function. Testosterone also plays a role in this regulation. The production of spermatozoa as well as the synthesis and release of testosterone are both the responsibility of the testes³². A lower testosterone level may be associated with a more complex condition, such as erectile dysfunction^{33,34}. Phosphodiesterase 5 (PDE-5) inhibitors, such as Sildenafil, Tadalafil, and others, are

typically prescribed to patients who have been diagnosed with erectile dysfunction. According to several studies, the effectiveness of PDE-5 inhibitors is improved when there is an adequate quantity of testosterone in the body³⁵. Erectile dysfunction may be caused by several causes, but the one that is caused by a lack of testosterone is likely treatable by using medicines that raise testosterone levels³³. Other causes of erectile dysfunction may not be so easily remedied. In the current study, after administering a treatment consisting of *C. nardus* extract for a period of 14 days, significant increases in testosterone levels were observed in the serum.

One of the herbal medications that have been shown in a recent study to have the potential impact of protecting organs against free radicals is called *C. nardus*^{36,19}. Antioxidants, flavonoids, phenolics, and terpenoids are the types of phytochemical substances that can be found in *C. nardus*^{37,38}. Antioxidants shield lipid membranes against damage caused by superoxide, ammonium, and peroxide radicals, which are each capable of causing cellular degeneration^{39,40}. Antioxidants also participate in the electron transport process that occurs during the final step of energy production in mitochondria⁴¹. To improve sperm motility, mitochondria must produce ATP or other forms of energy before fertilization may proceed⁴².

Fertilization is dependent on several factors, one of which is motility. If motility were to suffer any kind of impairment, this would have a negative influence on fertilization capacity⁴³. During their journey through the epididymis, sperm develop the ability to move about, and the epididymis is responsible for the initialization of this ability by both providing a distinct microenvironment and also secreting proteins that are essential for the initialization of sperm motility^{21,44}.

In the previous research, the protective effect of lemongrass in male reproduction after being triggered by hydrogen peroxide was described. And the results demonstrated that an increase in testosterone level, sperm characteristics, testicular and epididymal weight, as well as a reduction in serum and tissue homogenate *malondialdehyde* (MDA) and testicular histopathological changes, can be brought about by administering 100mg of lemongrass extract per kg of body weight⁴⁵. In correlation with the previous research, we found that *C. nardus* extract could improve sperm quality in mice after being induced by *S. aureus*. The potential dose of *C. nardus* that could improve sperm quality significantly was 25mg/kg of body weight, and this was found to be the optimal amount for this improvement.

CONCLUSION:

The *C. nardus* extract protects spermatozoa cells by repairing the sperm quality decrease caused by *S. aureus* infection. A low to a middle dose of *C. nardus* extract provides more protection than a high dose. A high dose of *C. nardus* extract is ineffective at improving sperm motility and concentration.

CONFLICT OF INTEREST:

The authors have no conflicts of interest regarding this research.

ACKNOWLEDGMENT:

We express our gratitude to the ministries of education, culture, research, and technology degree No. 2660/E4/AK.04/2021 for providing the grant for this research.

REFERENCES:

1. Borghat M Vander, Wyns C. Fertility and infertility: Definition and epidemiology. Clin Biochem. 2018; 62(March): 2-10.
2. Kaur S, Prabha V. Infertility as a Consequence of Spermagglutinating Staphylococcus aureus Colonization in Genital Tract of Female Mice. PLoS One. 2012;7(12). doi:10.1371/journal.pone.0052325
3. Avicenna F, Yudianto A, Itishom R, Wungu CDK. Effect of machine-washing semen-stained fabrics on the persistence of human spermatozoa DNA: A systematic review of five articles. Legal Medicine. 2023; 60: 102179. DOI:10.1016/j.legalmed.2022.102179
4. Wurlina W, Mustofa I, Meles DK, Mulyati S, Putri DKSC, Suwasanti N. Administration of the α -tocopherol for repairing testicle histological damage in rats exposed to dioxin. Thai Journal of Veterinary Medicine. 2021; 51(2): 293-301. DOI: 10.14456/tjvm.2021.37
5. Meles DK, Rachmawati K, Hamid IS, Mustofa I, Wurlina W, Suwasanti N, Putri DKSC, Utama S. α -Tocopherol Prevents Sperm Apoptosis and Necrosis in Rats Exposed to 2,3,7,8-Tetrachlorodibenzo-p-dioxin. Vet Med Int. 2022; 2022: 3685686. DOI: 10.1155/2022/3685686.
6. Esmailkhani A, Akhi MT, Sadeghi J, et al. Assessing the prevalence of Staphylococcus aureus in infertile male patients in Tabriz, northwest Iran. Int J Biomed. 2018;16(7):469-474.
7. Ghasemian F, Esmailnezhad S, Javad M, Moghaddam M. Staphylococcus saprophyticus and Escherichia coli: Tracking from sperm fertility potential to assisted reproductive outcomes. Clin Exp Reprod Med. 2021;48(2):142-149.
8. Sangavi R, Gopinath P, Kumar A. Antibacterial Activity of Ethanolic extract of Cinnamon against clinical Isolates of Staphylococcus aureus. Res J Pharm Technol. 2019;12(1):259-261.
9. Suryadinata, R. V., Sukarno, D. A., Sardjono, S. C., & Adriani, M. Antioxidant activity in red mulberries on sperm development exposed by cigarette smoke. Bali Medical Journal. 2021; 10(2): 583-586. DOI: 10.15562/bmj.v10i2.2329
10. Dutta S, Sengupta P, Izuka E, Menuba I, Jegasothy R. Staphylococcal infections and infertility: mechanisms and management. Mol Cell Biochem. 2020;474:57-72. doi:10.1007/s11010-020-03833-4
11. Sabudi IMNG. Factors influencing sperm retrieval rate by Microdissection Testicular Sperm Extraction (mTESE) in non-obstructive azoospermia cases: a literature review. Bali Medical Journal. 2020; 9(1), 356-359. doi: 10.15562/bmj.v9i1.1731
12. Darsini N, Hamidah B, Suyono SS, Ashari FY, Aswin RH, Yudiwati R. Human Sperm Motility, Viability, and Morphology

- Decreased after Cryopreservation. *Folia Medica Indonesiana*. 2019; 55(3): 198–201. DOI: 10.20473/fmi.v55i3.15501
13. Gopinath P. Antibacterial Activity of Ginger Oil against Clinical Isolates of *Staphylococcus aureus*. *Res J Pharm Technol*. 2018; 11(8): 3257-2=3258.
 14. Prameswara, I. G. N., Setiawan, A., Tanojo, T. D., Agustinus, A., & Hartono, J. Vitamin C and ubiquinone able to maintain sperm telomeres length on infertile's men: a clinical trial at Doctor Soetomo Regional General Hospital, Surabaya, Indonesia. *Bali Medical Journal*. 2020; 9(1): 360–365. DOI: 10.15562/bmj.v9i1.1733
 15. Sahu RK, Dewangan D, Roy A, Namdev KP, Vihar V. Anti-inflammatory Action of *Ougeinia oojenensis* (Roxb .) Hochr . Bark by HRBC Membrane Stabilization. *Res J Pharm Technol*. 2008;1(1):57-58.
 16. Guyansyah, A., Wratsangka, R., Dhanardono, D., Ghazali, M. F., Edy, H. J., Widyatama, H. G., Kusumaningrum, D., Tjahyadi, D., & Parwanto, E. Primary infertility of male and female factors, polycystic ovary syndrome and oligoasthenoteratozoospermia dominate the infertile population in agricultural and industrial areas in Karawang Regency, West Java Province, Indonesia. *Bali Medical Journal*. 2021; 10(1): 167–173. DOI: 10.15562/bmj.v10i1.2281
 17. Bagus Komang Satriyasa, I Gusti Ayu Widiyanti, & I.B.G. Fajar Manuaba. Depot Medroxyprogesterone acetate reduces spermatogonia cells and spermatid cells in the seminiferous tubules of male mice. *Bali Medical Journal*. 2022; 11(1): 508–512. DOI: 10.15562/bmj.v11i1.3459
 18. Abdullelah Z, Naqqash A, Al-bazaz HK, Salh M, Ibraheem SQ. GC-Mass and Phytochemical Investigation of *Cymbopogon citratus*. *Res J Pharm Technol*. 2019; 12(1): 67-73.
 19. Ayu P, Setiyowati I, Solekha R, Irawan DD, Rachmadani KA, Saputra AA. Protective Effect of Lemongrass Extract (*Cymbopogon nardus*) on Spermatozoa Cells of Lead Acetate Induced Mice. *Biomol Heal Sci J*. 2022; 05(01): 29-32. doi:10.20473/bhsj.v5i1.31450
 20. Kaur H, Bhardwaj U, Kaur R. *Cymbopogon nardus* essential oil : a comprehensive review on its chemistry and bioactivity. *J Essent Oil Res*. 2021; 33(3): 205-220. doi:10.1080/10412905.2021.1871976
 21. Zhang Y, Song XL, Yu B, Foong LC, Shu Y, Mai CW, Hu J, Dong B, Xue W, Chua CW. TP53 loss-of-function causes vulnerability to autophagy inhibition in aggressive prostate cancer. *Int J Urol*. 2022; 29(9): 1085-1094. doi: 10.1111/iju.15021.
 22. Akbari F, Azadbakht M, Gaurav A, Azimi F, Mahdizadeh Z, Vahedi L, Barzegar Nejad A, Chabra A, Eghbali M. Evaluation of the Therapeutic Effect of the Traditional Herbal Medicine *Atrifil* and *Oshagh Gum* on Testosterone-Induced Benign Prostatic Hyperplasia in Wistar Rats. *Adv Urol*. 2022; 2022: 5742431. DOI: 10.1155/2022/5742431
 23. Wang M, Wang Q, Du Y, Jiang H, Zhang X. Vitamins combined with traditional Chinese medicine for male infertility: A systematic review and meta-analysis. *Andrology*. 2020;8(5):1038-1050. doi:10.1111/andr.12787
 24. Kumar AS, Mazumder A, Vanitha J, Ganesh M, Venkateshwaran K, Saravanan VS. Antibacterial Activity of Methanolic Extract of *Sesbania Grandiflora* (Fabaceae). *Res J Pharm Technol*. 2008; 1(1): 59-60.
 25. C RC, Radhika J. Screening of Physicochemical properties and In Vitro free radical scavenging activity of *Delonix elata L .* leaf extracts. *Res J Pharm Technol*. 2020; 13(6): 2727-2732.
 26. Sinha A, Gupta S. Lipid Peroxidation and its Impact on Infertility. *Womens Heal Gynecol*. 2018; 4(1): 1-10.
 27. Wang S, Zhang K, Yao Y, Li J, Deng S. Bacterial Infections Affect Male Fertility: A Focus on the Oxidative Stress-Autophagy Axis. *Front Cell Dev Biol*. 2021; 9: 1-15. doi:10.3389/fcell.2021.727812
 28. Qasim MT, Al-mayali HK. Investigate the relation between Baicalin effect and Gene expression of LH , FSH , Testosterone in male rats treated with Gemcitabine drug. *Res J Pharm Technol*. 2018; 12(9): 4135-4141.
 29. Santi D, Cr P, Reiter E, et al. Follicle-Stimulating Hormone (FSH) Action on Spermatogenesis: A Focus on Physiological and Therapeutic Roles. *J Clin Med*. 2020; 9(4):1-27.
 30. Oduwale OO, Huhtaniemi IT, Misrahi M. The Roles of Luteinizing Hormone, Follicle-Stimulating Hormone and Testosterone in Spermatogenesis and Folliculogenesis Revisited. *Int J Mol Sci*. 2021;22(23):1-30.
 31. Susilowati S, Mustofa I, Wurlina W, Hernawati T, Oktanella Y, Soeharsono S, Purwanto DA. Green Tea Extract in the Extender Improved the Post-Thawed Semen Quality and Decreased Amino Acid Mutation of Kacang Buck Sperm. *Vet Sci*. 2022; 9(8): 403. doi:10.3390/vetsci9080403.
 32. Anaissie J, Delay KJ, Wang W, Hatzichristodoulou G, Hellstrom WJ. Testosterone deficiency in adults and corresponding treatment patterns across the globe. *Transl Androl Urol*. 2017;6(15):183-191. doi:10.21037/tau.2016.11.16
 33. Barkin J. Erectile dysfunction and hypogonadism (low testosterone). *Can J Urol*. 2011;18(April):2-7.
 34. Ibrahim MK, Tikamdas R, Kamal M. Testosterone Undecanoate effects on behavior and Cognitive Functions in male swiss Albino mice exposed to Chronic Social Defeat. *Res J Pharm Technol*. 2020;13(12):6041-6049.
 35. Pyrgidis N, Mykoniatis I, Haidich A, Bettina, Tirta M. The Effect of Phosphodiesterase-type 5 Inhibitors on Erectile Function: An Overview of Systematic Reviews. *Front Pharmacol*. 2021;12(September):1-11. doi:10.3389/fphar.2021.735708
 36. Saputra NA, Wibisono HS, Darmawan S, Pari G. Chemical composition of *Cymbopogon nardus* essential oil and its broad spectrum benefit. *IOP Conf Ser Earth Environ Sci*. 2020;415(1). doi:10.1088/1755-1315/415/1/012017
 37. Solekha, Rofiatun, Putri Ayu Ika Setiyowati SBSM, Kusumanegara CTUS. Phytochemical Screening of Ethanol Extract on Stems, Leaves, and Roots of Citronella Grass (*Cymbopogon nardus L.*). *BEST J (Biology Educ Sci Technol)*. 2022;5(1):141-147.
 38. Haque ANMA, Remadevi R, Naebe M. Lemongrass (*Cymbopogon*): a review on its structure, properties, applications and recent developments. *Cellulose*. 2018; 25(10): 5455-5477. doi:10.1007/s10570-018-1965-2
 39. Ma I, Ks C. Pharmacognostic Study of various species of *Cymbopogon* of South canara , India. *Res J Pharm Technol*. 2019;12(8):3626-3628.
 40. Abdulazeem L, Jassani MJAL, Al- MA. Free Radical Scavenging and Antioxidant Activity of Silver Nanoparticles Synthesized from *Cuminum cyminum* (Cumin) seed Extract. *Res J Pharm Technol*. 2021;14(8):4349-4356.
 41. Upadhyay PK, Dixit P, Garabadu D. Pharmacological Activities of Phytoconstituents and Essential oil obtained from *Cymbopogon citratus* Linn . *Res J Pharm Technol*. 2021;14(9).
 42. Prastiya RA, Suprayogi TW, Debora AE, Wijayanti A, Amalia A, Sulistyowati D, Nugroho AP. Green tea extract addition into a Tris-based egg yolk extender improves Bali bull sperm quality. *Anim Biosci*. 2023; 36(2): 209-217. DOI: 10.5713/ab.22.0184.
 43. Hakemi SG, Sc M, Shariffar F, et al. The Effects of Olive Leaf Extract on The Testis , Sperm Quality and Testicular Germ Cell Apoptosis i n Male Rats Exposed to Busulfan. *Int J Steril*. 2019;13(1):57-65. doi:10.22074/ijfs.2019.5520.Introduction
 44. Bertolla RP. Sperm biology and male reproductive health. *Sci Rep*. 2020;10:2-4. doi:10.1038/s41598-020-78861-7
 45. Rahim SM, Taha EM, Mubark ZM, Aziz SS, Simon KD, Mazlan AG. Protective effect of *Cymbopogon citratus* on hydrogen peroxide-induced oxidative stress in the reproductive system of male rats. *Syst Biol Reprod Med*. 2013;59(6):329-336. doi:10.3109/19396368.2013.827268

64_RJPT_17_2

by Putri Setiyowati

Submission date: 26-Feb-2024 05:38AM (UTC-0600)

Submission ID: 2304891168

File name: 64_RJPT_17_2_2024.pdf (306.71K)

Word count: 4920

Character count: 26297

RESEARCH ARTICLE

Examining the effects of Lemongrass (*Cymbopogon nardus*) extract on Sperm Quality and Testosterone Level in Mice Induced with *Staphylococcus aureus*

Lilis Maghfuroh¹, Putri Ayulka Setiyowati^{2*}, Rofiatun Solekha²,
Alif'iyah Mar' Atus Sholikhah¹, Angella Ananda Saputra², Eka Febrianti Wulandari¹

¹Nursing Program, Faculty of Health Science, Universitas Muhammadiyah Lamongan, East Java, Indonesia.

²Biology Program, Faculty of Science, Technology, and Education,
Universitas Muhammadiyah Lamongan, East Java, Indonesia.

*Corresponding Author E-mail: putriayuikasetiyowati@gmail.com

ABSTRACT:

There are a variety of causes for male infertility. Among them is microbial infection. Lemongrass (*C. nardus*) contains natural antioxidants and flavonoids that have the ability to preserve the male reproductive system. This study examined the effects of different *C. nardus* extract dosages on mouse sperm quality. The male Balb/C mice were separated into five groups of equal size. The negative control group was injected with 0.01ml of Na-CMC 0.5% over the course of 28 days, the positive control group was injected with 0.2ml of *Staphylococcus aureus* (*S. aureus*) on days 10, 17, and 24, and the treatment groups were injected with 0.2ml of *S. aureus* on days 10, 17, and 24, followed by 0.01ml of *C. nardus*. After the experiment, mice were slaughtered, and sperm suspension was obtained from the cauda epididymal to measure morphology, concentration, and motility. The testis was fixed in a fixative solution to analyze the spermatogenic cells, and intracardiac blood was drawn to determine the level of testosterone. Significant differences ($p < 0.05$) were seen in the morphology, concentration, and motility of spermatozoa when *C. nardus* extract was administered. Given the low and middle concentrations of *C. nardus* extract, it can repair sperm damage caused by *S. aureus* infection.

KEYWORDS: Male infertility, *C. nardus* extract, *S. aureus*, Sperm quality, Testosterone level.

INTRODUCTION:

Infertility is becoming more common in Indonesia every year. Infertility affects not only women but also men. Men are responsible for 36% of fertility success^{1,2}. Male infertility can be caused by a variety of factors, including bacteria pathogens such as *Chlamydia sp.*, *Escherichia coli*, *Mycoplasma hominis*, and *Staphylococcus aureus*. This bacteria has been shown to reduce sperm motility^{3,4,5}. *Staphylococcus aureus* (*S. aureus*) can cause infections in a variety of ecological niches within the host⁶. It colonizes the nares, axillae, vagina, pharynx, and damaged skin surfaces in humans, causing a variety of suppurative (pus-forming) infections and toxins^{7,8}.

Many studies have revealed that *S. aureus* can cause male infertility^{9,10}. One study revealed that incubating *S. aureus* in sperm reduced testis weight, seminiferous tubulous diameter, epididymal sperm count, and serum testosterone level^{11,12}. Another study found that *S. aureus* alters the composition of sperm proteins and promotes early cell death by secreting lipopolysaccharides (LPS). It resulted in an increase in dead cells and a decrease in motile sperm cells^{3,13}. In this case, the use of antioxidant-containing plants is thought to reduce the toxic effect of various toxicants, including bacterial infection^{14,15}. The concentration of spermatozoa in one ejaculation depends on the process of formation of spermatogenesis in seminiferous tubules¹⁶. When spermatogenesis proceeds normally it will be generated a normal spermatozoa count, otherwise during the process spermatogenesis occurs interference, then the development of spermatogonia cells will affect the number of spermatozoa formed. It

Received on 28.11.2022 Modified on 15.02.2023
Accepted on 03.04.2023 © RJPT All right reserved
Research J. Pharm. and Tech 2024; 17(2):879-884.
DOI: 10.52711/0974-360X.2024.00136

depends on the magnitude of the disturbance that occurred during the process of spermatogenesis^{16,17}. Lemongrass is a natural antioxidant with numerous benefits that are commonly used in herbal medicine¹⁸. *Cymbopogon nardus*, one type of lemongrass studied in this study, is known for its antioxidant activity and effectiveness to attack bacteria by lysis the cells via its potent flavonoid compound^{19,20}. As a result, this study looked into the potential protective effects of *C. nardus* extract on the male reproductive system after infection with *S. aureus*. The parameters measured were epididymal and testicular weight, the morphology of sperm, concentration of sperm, motility of sperm, spermatogenic cells, the diameter of seminiferous tubules, and testosterone level.

MATERIALS AND METHODS:

Ethical approval:

With the number 120-KEP-UB-2021, the Ethical Review Committee of the Research and Community Service Department at Universitas Brawijaya, East Java, Indonesia, approved all procedures in this study, including the use of mice as animal models.

Plant-based substances:

C. nardus was collected in September 2021 from Materia Medica, Batu, East Java. It was acknowledged and authenticated by Universitas Muhammadiyah Lamongan's Biological Laboratory, Biology Department.

Na-CMC 0.5% preparation and suspension:

A total of 500 milligrams of *Na-Carboxymethyl cellulose* (Na-CMC) were weighed and dissolved in warm distilled water. Using a 100ml measuring flask, all of the remaining distilled water was added after dissolving to obtain a volume of 100ml Na-CMC solution.

C. nardus ethanolic extract preparation and suspension:

C. nardus stalks were cut into little pieces and dried in the sun for about five days. The 250g of dried stalks were ground in an electric grinder before being macerated with ethanol at room temperature for three days. Then, the extract was filtered and concentrated in a water bath at 70⁰ degrees Celsius. To create the suspension, the extract was homogeneously mashed with 0.5% Na-CMC. The respective concentrations were 0.5%, 1%, and 2%.

Microorganism:

The strain of *S. aureus* employed in this investigation was previously obtained from the cervix of an infertile woman in our laboratory.

Substances and chemicals:

Ethanol, methanol, eosin, nigrosine, chloroform (purchased from Merck, Darmstadt, Germany), distilled water, formalin 10% neutral buffer, paraffin, hematoxyline eosin, testosterone kit, xylol, alcohol 70%, alcohol 96%, and NaCl 0.9%.

Animals:

Thirty mature male Balb/C mice were donated from the Veterinary and Pharma Center in Surabaya, East Java (eight weeks old, weighing 25–30grams). They were fed and watered ad libitum in standard laboratory conditions (temperature room at 28–30°C, and 12-h–12-h light-dark cycle).

Experimental design:

After a week of acclimatization, the animals were put into five equal groups (n = 6). The negative control groups were given 0.1ml of Na-CMC 0.5% subcutaneously every day for 28 days. The positive control groups were given 0.2ml of *S. aureus* (0.5 McFarland) three times on days 10, 17, and 24. As positive controls, 0.2ml *S. aureus* (0.5McFarland) was given three times to the treatment groups through the intraperitoneal route. Then, different doses of 0.1ml were given subcutaneously to each treatment group in a steady stream. The first group was given 25mg/kg of body weight, the second group was given 50mg/kg of body weight, and the third group was given 100mg/kg of body weight. All three treatments were given over the duration of 28 days. At the end of the process, chloroform was used to kill all of the mice. Then, the cauda epididymal was taken and used to make a sperm suspension for measuring sperm.

Sperm analysis:

This study initially measured epididymal and testicular weight from each sample, then measured the percentage of motility, morphology, and the concentration of spermatozoa in a suspension. 10µL was evaluated under a light microscope in a hemocytometer chamber for sperm motility. One hundred sperm cells were counted in each animal, and the percentage of motility was classified as motil (%) and immotile (%). The motile sperms have moved straight. An hour after sperm diffusion in the solution, 10µL of the sperm suspension was transferred to each hemocytometer counting chamber for 5 minutes to evaluate sperm concentration. Using a 40-magnification light microscope, sperm heads were counted and quantified as million/ml of suspension. Eosin-nigrosine staining determined sperm morphology. To make smears, spermatozoa suspension is dropped on the object's glass and dried in the air. Smears were fixed with methanol, colored with 1% Y eosin solution and nigrosine, and dried. Distilled water cleaned and dried the preparations. 100 mouse

spermatozoa were examined using a 400-magnification light microscope. Calculating the normal/abnormal spermatozoa percentage was the final stage.

Levels of Testosterone:

Serum was taken from intracardiac blood. An enzyme-linked immunoassay kit (Bioassay Technology Lab® 96 kit testosterone, China) evaluated serum testosterone. The level of testosterone measured as nmol/mL.

Examining Spermatogenic Cells Histologically:

After fixing the testicular collection in neutral buffer formalin 10%, paraffin, and hematoxyline eosin were used to make histological preparations with a 5µm thickness. The diameter of seminiferous tubules and the number of spermatogenic cells (spermatogonia, spermatid, and spermatocytes) were examined in mouse testicular preparations under 400x magnification.

Data analysis:

All data were represented as means standard error of the mean (SEM), and statistical analysis was conducted using one-way analysis of variance (one-way ANOVA) and post hoc multiple comparison test to determine differences between pairs of means.

RESULT:

The results showed that giving *S. aureus* to mice for 28 days led to a significant ($p < 0.05$) decrease in the quantity of sperm and the percentage of spermatozoa that could move (immotile). In mice that were infected by *S. aureus*, the percentage of spermatozoa that had an aberrant morphology was statistically ($p < 0.05$) higher than it was in the group that served as a negative control (K1). When compared to animals infected with *S. aureus*, testosterone levels and sperm parameters such as motility and concentration increased significantly ($p < 0.05$) in mice that were given *C. nardus* extract (Table 1). The diameter of the seminiferous tubule, the quantity of spermatogonia cells, spermatocytes, and spermatid

cells in the positive control that was infected by *S. aureus* all had lower values than the normal negative control and the group that was treated with *C. nardus* extract. This was another finding (Table 2).

According to the data in table 1, spermatozoa motility at a dose of 100 mg/kg body weight of *C. nardus* extract (P3) was not significantly different from the positive control group (K2). This means that at high doses, the ability to restore the quality of spermatozoa damaged by lead acetate exposure was not optimal. Furthermore, at a dose of 50 mg/kg body weight, the *C. nardus* extract demonstrated the highest average in each observational parameter. It is reasonable to assume that the dose was optimal for improving spermatozoa quality. A description of sperm morphological observation was shown in Figure 1. The differences in the histological seminiferous tubule in each group were shown in figure 2.




Fig.1: Sperm cell morphology in adult mice had normal (N) and abnormal (A: twisted midpiece or tail; B: head malformation (like a needle); C: larger head; D: broken tail).

Table 1: The effect of *C. nardus* extract on sperm motility, morphology, concentration, and testosterone levels in *S. aureus*-infected mice.

Parameters	K1	K2	P1	P2	P3
Motility (%)	80.60 ± 2.51 ^a	58.20 ± 2.28 ^b	70.40 ± 2.70 ^{ab}	76.20 ± 4.02 ^a	61.00 ± 1.58 ^b
Normal morphology (%)	83.60 ± 2.30 ^a	41.60 ± 2.07 ^{bc}	67.20 ± 1.92 ^{ab}	70.40 ± 3.36 ^{ab}	51.00 ± 2.92 ^b
Concentration (x10 ⁶ /mL)	83.00 ± 2.55 ^a	50.00 ± 2.24 ^{ab}	72.20 ± 1.48 ^b	77.20 ± 1.92 ^{bc}	65.80 ± 2.78 ^c
Testosterone level (nmol/L)	13.68 ± 0.65 ^a	8.60 ± 0.40	10.90 ± 0.57	12.89 ± 0.85 ^a	9.69 ± 0.42
Testicular weight (mg)	306 ± 20.73 ^a	216 ± 20.73 ^{ab}	258 ± 23.76 ^b	292 ± 23.88 ^a	245 ± 7.07 ^{ab,b}
Epididymal weight (mg)	10.84 ± 0.39 ^a	7.68 ± 0.40	10.02 ± 0.58 ^{ab}	10.36 ± 0.63 ^{ab}	8.46 ± 0.34

Data are presented as mean ± SD. Differences in superscripts in the same rows show significant differences ($p < 0.05$).

K1 (negative control): Na-CMC 0.5%; K2 (positive control): *S. aureus* 0.5 Mc Farland; P1 (treatment group 1): *S. aureus* 0.5 Mc Farland + *C. nardus* 25 mg/kg BW; P2 (treatment group 2): *S. aureus* 0.5 Mc Farland + *C. nardus* 50 mg/kg BW; P3 (treatment group 3): *S. aureus* 0.5 Mc Farland + *C. nardus* 100 mg/kg BW.




Fig. 2: Testicular seminiferous tubules in different groups. Red arrows indicate spermatogonia cells, yellow spermatocytes, blue spermatids, and black seminiferous tubule diameter. (Hematoxylin Eosin: 400x).

Table 2. Effect of *C. nardus* extract on spermatogenic cells and diameter of the seminiferous tubule in mice infected by *S. aureus*

Parameters	K1	K2	P1	P2	P3
DST	216.52 ± 16.65 ^a	105.60 ± 13.91 ^b	141.41 ± 13.19	186.03 ± 29.70 ^a	115.79 ± 13.85 ^b
Spermatogonia cells	82.80 ± 4.38 ^a	60.00 ± 4.00	75.20 ± 4.60 ^{ab}	80.00 ± 3.16 ^{a,ab}	67.20 ± 4.15
Spermatocytes cells	99.60 ± 5.73 ^a	65.20 ± 7.95	87.80 ± 6.72 ^{ab}	98.00 ± 6.33 ^a	81.20 ± 3.35 ^{ab}
Spermatid cells	284.0 ± 42.9 ^a	179.2 ± 18.6	255.2 ± 39.5 ^{a,ab}	268.0 ± 24.5 ^{a,ab}	204.8 ± 15.1

Data are presented as mean ± SD

Differences in superscripts in the same rows show significant differences ($p < 0.05$). K1 (negative control): Na-CMC 0.5%; K2 (positive control): *S. aureus* 0.5 Mc Farland; P1 (treatment group 1): *S. aureus* 0.5 Mc Farland + *C. nardus* 25 mg/kg BW; P2 (treatment group 2): *S. aureus* 0.5 Mc Farland + *C. nardus* 50 mg/kg BW; P3 (treatment group 3): *S. aureus* 0.5 Mc Farland + *C. nardus* 100 mg/kg BW. DST= Diameter of Seminiferous Tubules

DISCUSSION:

In ethnomedicines, the use of medicinal plants, commonly known as medicinal herbs, has been suggested for the treatment of a wide variety of disorders at various points in history and continued to be practiced today. Throughout the history of humanity, people have relied on the therapeutic properties of various plants to heal a variety of illnesses^{21,22}. New scientific approaches have been integrated into conventional medical practice to facilitate the standardization of phytotherapy, which refers to the use of numerous plant-based remedies in combination with one another^{23,14}.

According to this study, The decreased quality of spermatozoa that has been caused by bacterial infection was partially restored by the use of an extract of *C. nardus*. There was a statistically significant difference

($p < 0.05$) between the group that was treated with *C. nardus* extract and the group that served as the positive control and was treated with *S. aureus*. To acquire the optimal results in terms of epididymal and testicular weight, sperm motility, normal morphology of sperm, testosterone level, sperm concentration, and the number of spermatogenic cells, a dose of 25mg/kg of body weight was shown to be the most effective. The increased dose that was administered did not result in an improvement in the quality of the spermatozoa. Even at a dose of 100mg/kg of body weight, the quality of sperm that had been infected with *S. aureus* was not improved.

S. aureus is a pathogen bacteria that are typically found in the female genital tract². This bacteria is known to have a toxic effect on the epididymal duct in addition to disrupting the process of spermatogenesis and also could well interact with the CD14 on the surface of the sperm, primarily in the head and tail region²⁴. This interaction will result in increased levels of sperm membrane lipid peroxidation, production of reactive oxygen species, and caspase-mediated apoptosis in the subsequent stages^{6,10,25}. A reduction in the motility of spermatozoa and an increase in lipid peroxidation are both indicators that cell damage has been caused by *S. aureus*^{26,16,27}.

The process of male reproduction is a complicated one that requires a great deal of energy. The hypothalamus and the pituitary gland use a variety of feedback regulation mechanisms to control the function of the testis, which serves as the primary organ of the male germline^{1,28}. GnRH, FSH, and LH are the primary hormones responsible for controlling regulatory mechanisms via their actions on the hypothalamus-pituitary-testis axis²⁹. Through neurons that secrete GnRH, the hypothalamus can monitor the condition of the reproductive system. Through the anterior pituitary, GnRH stimulates the release of LH as well as FSH. LH stimulates the production of testosterone by acting on testicular interstitial cells and increasing their permeability. FSH is the hormone that is responsible for the stimulation of the seminiferous tubules, which is necessary to maintain sperm production³⁰.

A decrease in the amount of testosterone in the body can lead to reproductive disorders. Testosterone is a hormone that is primarily found in men³¹. The precise and coordinated secretion of GnRH, LH, FSH, and testosterone is what regulates male reproductive function. Testosterone also plays a role in this regulation. The production of spermatozoa as well as the synthesis and release of testosterone are both the responsibility of the testes³². A lower testosterone level may be associated with a more complex condition, such as erectile dysfunction^{33,34}. Phosphodiesterase 5 (PDE-5) inhibitors, such as Sildenafil, Tadalafil, and others, are

typically prescribed to patients who have been diagnosed with erectile dysfunction. According to several studies, the effectiveness of PDE-5 inhibitors is improved when there is an adequate quantity of testosterone in the body³⁵. Erectile dysfunction may be caused by several causes, but the one that is caused by a lack of testosterone is likely treatable by using medicines that raise testosterone levels³³. Other causes of erectile dysfunction may not be so easily remedied. In the current study, after administering a treatment consisting of *C. nardus* extract for a period of 14 days, significant increases in testosterone levels were observed in the serum.

One of the herbal medications that have been shown in a recent study to have the potential impact of protecting organs against free radicals is called *C. nardus*^{36,19}. Antioxidants, flavonoids, phenolics, and terpenoids are the types of phytochemical substances that can be found in *C. nardus*^{37,38}. Antioxidants shield lipid membranes against damage caused by superoxide, ammonium, and peroxide radicals, which are each capable of causing cellular degeneration^{39,40}. Antioxidants also participate in the electron transport process that occurs during the final step of energy production in mitochondria⁴¹. To improve sperm motility, mitochondria must produce ATP or other forms of energy before fertilization may proceed⁴².

Fertilization is dependent on several factors, one of which is motility. If motility were to suffer any kind of impairment, this would have a negative influence on fertilization capacity⁴³. During their journey through the epididymis, sperm develop the ability to move about, and the epididymis is responsible for the initialization of this ability by both providing a distinct microenvironment and also secreting proteins that are essential for the initialization of sperm motility^{21,44}.

In the previous research, the protective effect of lemongrass in male reproduction after being triggered by hydrogen peroxide was described. And the results demonstrated that an increase in testosterone level, sperm characteristics, testicular and epididymal weight, as well as a reduction in serum and tissue homogenate *malondialdehyde* (MDA) and testicular histopathological changes, can be brought about by administering 100mg of lemongrass extract per kg of body weight⁴⁵. In correlation with the previous research, we found that *C. nardus* extract could improve sperm quality in mice after being induced by *S. aureus*. The potential dose of *C. nardus* that could improve sperm quality significantly was 25mg/kg of body weight, and this was found to be the optimal amount for this improvement.

CONCLUSION:

The *C. nardus* extract protects spermatozoa cells by repairing the sperm quality decrease caused by *S. aureus* infection. A low to a middle dose of *C. nardus* extract provides more protection than a high dose. A high dose of *C. nardus* extract is ineffective at improving sperm motility and concentration.

CONFLICT OF INTEREST:

The authors have no conflicts of interest regarding this research.

ACKNOWLEDGMENT:

We express our gratitude to the ministries of education, culture, research, and technology degree No. 2660/E4/AK.04/2021 for providing the grant for this research.

REFERENCES:

1. Borghet M Vander, Wyns C. Fertility and infertility: Definition and epidemiology. Clin Biochem. 2018; 62(March): 2-10.
2. Kaur S, Prabha V. Infertility as a Consequence of Spermagglutinating Staphylococcus aureus Colonization in Genital Tract of Female Mice. PLoS One. 2012;7(12). doi:10.1371/journal.pone.0052325
3. Avicenna F, Yudianto A, Itishom R, Wungu CDK. Effect of machine-washing semen-stained fabrics on the persistence of human spermatozoa DNA: A systematic review of five articles. Legal Medicine. 2023; 60: 102179. DOI:10.1016/j.legalmed.2022.102179
4. Wurlina W, Mustofa I, Meles DK, Mulyati S, Putri DKSC, Suwasanti N. Administration of the α -tocopherol for repairing testicle histological damage in rats exposed to dioxin. Thai Journal of Veterinary Medicine. 2021; 51(2): 293-301. DOI: 10.14456/tjvm.2021.37
5. Meles DK, Rachmawati K, Hamid IS, Mustofa I, Wurlina W, Suwasanti N, Putri DKSC, Utama S. α -Tocopherol Prevents Sperm Apoptosis and Necrosis in Rats Exposed to 2,3,7,8-Tetrachlorodibenzo-p-dioxin. Vet Med Int. 2022; 2022: 3685686. DOI: 10.1155/2022/3685686.
6. Esmailkhani A, Akhi MT, Sadeghi J, et al. Assessing the prevalence of Staphylococcus aureus in infertile male patients in Tabriz, northwest Iran. Int J Biomed. 2018;16(7):469-474.
7. Ghasebian F, Esmailnezhad S, Javad M, Moghaddam M. Staphylococcus saprophyticus and Escherichia coli: Tracking from sperm fertility potential to assisted reproductive outcomes. Clin Exp Reprod Med. 2021;48(2):142-149.
8. Sangavi R, Gopinath P, Kumar A. Antibacterial Activity of Ethanolic extract of Cinnamon against clinical Isolates of Staphylococcus aureus. Res J Pharm Technol. 2019;12(1):259-261.
9. Suryadinata, R. V., Sukarno, D. A., Sardjono, S. C., & Adriani, M. Antioxidant activity in red mulberries on sperm development exposed by cigarette smoke. Bali Medical Journal. 2021; 10(2): 583-586. DOI: 10.15562/bmj.v10i2.2329
10. Dutta S, Sengupta P, Izuka E, Menuba I, Jegasothy R. Staphylococcal infections and infertility: mechanisms and management. Mol Cell Biochem. 2020;474:57-72. doi:10.1007/s11010-020-03833-4
11. Sabudi IMG. Factors influencing sperm retrieval rate by Microdissection Testicular Sperm Extraction (mTESE) in non-obstructive azoospermia cases: a literature review. Bali Medical Journal. 2020; 9(1), 356-359. doi: 10.15562/bmj.v9i1.1731
12. Darsini N, Hamidah B, Suyono SS, Ashari FY, Aswin RH, Yudiwati R. Human Sperm Motility, Viability, and Morphology

- Decreased after Cryopreservation. *Folia Medica Indonesiana*. 2019; 55(3): 198–201. DOI: 10.20473/fmi.v55i3.15501
13. Gopinath P. Antibacterial Activity of Ginger Oil against Clinical Isolates of *Staphylococcus aureus*. *Res J Pharm Technol*. 2018; 11(8): 3257-2=3258.
14. Pramesemara, I. G. N., Setiawan, A., Tanojo, T. D., Agustinus, A., & Hartono, J. Vitamin C and ubiquinone able to maintain sperm telomeres length on infertile's men: a clinical trial at Doctor Soetomo Regional General Hospital, Surabaya, Indonesia. *Bali Medical Journal*. 2020; 9(1): 360–365. DOI: 10.15562/bmj.v9i1.1733
15. Sahu RK, Dewangan D, Roy A, Namdev KP, Vihar V. Anti-inflammatory Action of *Ougeinia oojensis* (Roxb .) Hochr . Bark by HRBC Membrane Stabilization. *Res J Pharm Technol*. 2008;1(1):57-58.
16. Guyansyah, A., Wratsangka, R., Dhanardono, D., Ghazali, M. F., Edy, H. J., Widyatama, H. G., Kusumaningrum, D., Tjahyadi, D., & Parwanto, E. Primary infertility of male and female factors, polycystic ovary syndrome and oligoasthenoteratozoospermia dominate the infertile population in agricultural and industrial areas in Karawang Regency, West Java Province, Indonesia. *Bali Medical Journal*. 2021; 10(1): 167–173. DOI: 10.15562/bmj.v10i1.2281
17. Bagus Komang Satriyasa, I Gusti Ayu Widiati, & I.B.G. Fajar Manuaba. Depot Medroxyprogesterone acetate reduces spermatogonia cells and spermatid cells in the seminiferous tubules of male mice. *Bali Medical Journal*. 2022; 11(1): 508–512. DOI: 10.15562/bmj.v11i1.3459
18. Abdullelah Z, Naqash A, Al-bazaz HK, Salh M, Ibraheem SQ. GC-Mass and Phytochemical Investigation of *Cymbopogon citratus*. *Res J Pharm Technol*. 2019; 12(1): 67-73.
19. Ayu P, Setiyowati I, Solekha R, Irawan DD, Rachmadani KA, Saputra AA. Protective Effect of Lemongrass Extract (*Cymbopogon nardus*) on Spermatozoa Cells of Lead Acetate Induced Mice. *Biomol Heal Sci J*. 2022; 05(01): 29-32. doi:10.20473/bhsj.v5i1.31450
20. Kaur H, Bhardwaj U, Kaur R. *Cymbopogon nardus* essential oil : a comprehensive review on its chemistry and bioactivity. *J Essent Oil Res*. 2021; 33(3): 205-220. doi:10.1080/10412905.2021.1871976
21. Zhang Y, Song XL, Yu B, Foong LC, Shu Y, Mai CW, Hu J, Dong B, Xue W, Chua CW. TP53 loss-of-function causes vulnerability to autophagy inhibition in aggressive prostate cancer. *Int J Urol*. 2022; 29(9): 1085-1094. doi: 10.1111/iju.15021.
22. Akbari F, Azadbakht M, Gaurav A, Azimi F, Mahdizadeh Z, Vahedi L, Barzegar Nejad A, Chabra A, Eghbali M. Evaluation of the Therapeutic Effect of the Traditional Herbal Medicine *Atrifol* and *Oshagh Gum* on Testosterone-Induced Benign Prostatic Hyperplasia in Wistar Rats. *Adv Urol*. 2022; 5742431. DOI: 10.1155/2022/5742431
23. Wang M, Wang Q, Du Y, Jiang H, Zhang X. Vitamins combined with traditional Chinese medicine for male infertility: A systematic review and meta-analysis. *Andrology*. 2020;8(5):1038-1050. doi:10.1111/andr.12787
24. Kumar AS, Mazumder A, Vanitha J, Ganesh M, Venkateshwaran K, Saravanan VS. Antibacterial Activity of Methanolic Extract of *Sesbania Grandiflora* (Fabaceae). *Res J Pharm Technol*. 2008; 1(1): 59-60.
25. C RC, Radhika J. Screening of Physicochemical properties and In Vitro free radical scavenging activity of *Delonix elata* L . leaf extracts. *Res J Pharm Technol*. 2020; 13(6): 2727-2732.
26. Sinha A, Gupta S. Lipid Peroxidation and its Impact on Infertility. *Womens Heal Gynecol*. 2018; 4(1): 1-10. 7
27. Wang S, Zhang K, Yao Y, Li J, Deng S. Bacterial Infections Affect Male Fertility: A Focus on the Oxidative Stress-Autophagy Axis. *Front Cell Dev Biol*. 2021; 9: 1-15. doi:10.3389/fcell.2021.727812 12
28. Qasim MT, Al-mayali HK. Investigate the relation between Baicalin effect and Gene expression of LH , FSH , Testosterone in male rats treated with Gemcitabine drug. *Res J Pharm Technol*. 2018; 12(9): 4135-4141.
29. Santi D, Cr P, Reiter E, et al. Follicle-Stimulating Hormone (FSH) Action on Spermatogenesis: A Focus on Physiological and Therapeutic Roles. *J Clin Med*. 2020; 9(4):1-27.
30. Oduwale OO, Huhtaniemi IT, Misrahi M. The Roles of Luteinizing Hormone, Follicle-Stimulating Hormone and Testosterone in Spermatogenesis and Folliculogenesis Revisited. *Int J Mol Sci*. 2021;22(23):1-30.
31. Susilowati S, Mustofa I, Wurlina W, Hernawati T, Oktanella Y, Soeharsono S, Purwanto DA. Green Tea Extract in the Extender Improved the Post-Thawed Semen Quality and Decreased Amino Acid Mutation of Kacang Buck Sperm. *Vet Sci*. 2022; 9(8): 403. doi: 10.3390/vetsci9080403.
32. Anaissie J, Delay KJ, Wang W, Hatzichristodoulou G, Hellstrom WJ. Testosterone deficiency in adults and corresponding treatment patterns across the globe. *Transl Androl Urol*. 2017;6(15):183-191. doi:10.21037/tau.2016.11.16
33. Barkin J. Erectile dysfunction and hypogonadism (low testosterone). *Can J Urol*. 2011;18(April):2-7.
34. Ibrahim MK, Tikamdas R, Kamal M. Testosterone Undecanoate effects on behavior and Cognitive Functions in male swiss Albino mice exposed to Chronic Social Defeat. *Res J Pharm Technol*. 2020;13(12):6041-6049.
35. Pyrgidis N, Mykoniatis I, Haidich A, Bettina, Tirta M. The Effect of Phosphodiesterase-type 5 Inhibitors on Erectile Function: An Overview of Systematic Reviews. *Front Pharmacol*. 2021;12(September):1-11. doi:10.3389/fphar.2021.735708
36. Saputra NA, Wibisono HS, Darmawan S, Pari G. Chemical composition of *Cymbopogon nardus* essential oil and its broad spectrum benefit. *IOP Conf Ser Earth Environ Sci*. 2020;415(1). doi:10.1088/1755-1315/415/1/012017
37. Solekha, Rofiatun, Putri Ayu Ika Setiyowati SBSM, Kusumanegara CTUS. Phytochemical Screening of Ethanol Extract on Stems, Leaves, and Roots of Citronella Grass (*Cymbopogon nardus* L.). *BEST J (Biology Educ Sci Technol*. 2022;5(1):141-147.
38. Haque ANMA, Remadevi R, Naebe M. Lemongrass (*Cymbopogon*): a review on its structure, properties, applications and recent developments. *Cellulose*. 2018; 25(10): 5455-5477. doi:10.1007/s10570-018-1965-2
39. Ma I, Ks C. Pharmacognostic Study of various species of *Cymbopogon* of South canara , India. *Res J Pharm Technol*. 2019;12(8):3626-3628.
40. Abdulazeem L, Jassani MJAL, Al- MA. Free Radical Scavenging and Antioxidant Activity of Silver Nanoparticles Synthesized from *Cuminum cyminum* (Cumin) seed Extract. *Res J Pharm Technol*. 2021;14(8):4349-4356.
41. Upadhyay PK, Dixit P, Garabadu D. Pharmacological Activities of Phytoconstituents and Essential oil obtained from *Cymbopogon citratus* Linn . *Res J Pharm Technol*. 2021;14(9).
42. Prastiya RA, Suprayogi TW, Debora AE, Wijayanti A, Amalia A, Sulistyowati D, Nugroho AP. Green tea extract addition into a Tris-based egg yolk extender improves Bali bull sperm quality. *Anim Biosci*. 2023; 36(2): 209-217. DOI: 10.5713/ab.22.0184.
43. Hakemi SG, Sc M, Shariffar F, et al. The Effects of Olive Leaf Extract on The Testis , Sperm Quality and Testicular Germ Cell Apoptosis i n Male Rats Exposed to Busulfan. *Int J Steril*. 2019;13(1):57-65. doi:10.22074/ijfs.2019.5520.Introduction
44. Bertolla RP. Sperm biology and male reproductive health. *Sci Rep*. 2020;10:2-4. doi:10.1038/s41598-020-78861-7
45. Rahim SM, Taha EM, Mubark ZM, Aziz SS, Simon KD, Mazlan AG. Protective effect of *Cymbopogon citratus* on hydrogen peroxide-induced oxidative stress in the reproductive system of male rats. *Syst Biol Reprod Med*. 2013;59(6):329-336. doi:10.3109/19396368.2013.827268

ORIGINALITY REPORT

11 %
SIMILARITY INDEX

0 %
INTERNET SOURCES

11 %
PUBLICATIONS

0 %
STUDENT PAPERS

PRIMARY SOURCES

- 1

Erfan Ghaderian, Bahman Rahimi Esboei, Parisa Mousavi, Maryam Pourhajibagher, Mohammad Mohsen Homayouni, Mohammad Zeinali. "Anti-leishmanial effects of Eryngium planum and Ecbillium elaterum methanolic extract against Leishmania major", AMB Express, 2024
Publication

1 %
- 2

James Anaissie, Kent J. DeLay, William Wang, Georgios Hatzichristodoulou, Wayne J. Hellstrom. "Testosterone deficiency in adults and corresponding treatment patterns across the globe", Translational Andrology and Urology, 2017
Publication

1 %
- 3

Xiao-Ying Liu, Yan-Ling Deng, Pan-Pan Chen, Chong Liu et al. "Self-Rated Health and Semen Quality in Men Undergoing Assisted Reproductive Technology", JAMA Network Open, 2024
Publication

1 %

4

Bin Wang, Jihua Xu, Shenhui Jiang, Yanbo Wang, Jun Zhu, Yugang Zhang. "Combined Analysis of Gut Microbiota and Plasma Metabolites Reveals the Effect of Red-Fleshed Apple Anthocyanin Extract on Dysfunction of Mice Reproductive System Induced by Busulfan", *Frontiers in Nutrition*, 2022

Publication

1 %

5

André Rizzuti, Conrado Alvarenga, Gustavo Stocker, Lucas Fraga, Heitor O. Santos. "Early Pharmacologic Approaches to Avert Anabolic Steroid-induced Male Infertility: A Narrative Review", *Clinical Therapeutics*, 2023

Publication

1 %

6

Marziye Farsimadan, Mohammad Motamedifar. "Bacterial infection of the male reproductive system causing infertility", *Journal of Reproductive Immunology*, 2020

Publication

1 %

7

Bushra A. Rakha, Zartasha Zafar, Muhammad S. Ansari, Shamim Akhter et al. "Influence of Bacterial Contamination and Antibiotic Sensitivity on Cryopreserved Sperm Quality of Indian Red Jungle Fowl", *Biopreservation and Biobanking*, 2023

Publication

<1 %

8

Divan Coetzee, Jiri Militky, Mohanapriya Venkataraman. "Functional Coatings by

<1 %

Natural and Synthetic Agents for Insect Control and Their Applications", Coatings, 2022

Publication

9

Ioannis Mykoniatis, Nikolaos Pyrgidis, Filimon Zilotis, Paraskeui Kapoteli et al. "The Effect of Combination Treatment With Low-Intensity Shockwave Therapy and Tadalafil on Mild and Mild-To-Moderate Erectile Dysfunction: A Double-Blind, Randomized, Placebo-Controlled Clinical Trial", The Journal of Sexual Medicine, 2022

Publication

<1 %

10

Ravindran Jegasothy, Pallav Sengupta, Sulagna Dutta, Ravichandran Jeganathan. "Climate change and declining fertility rate in Malaysia: the possible connexions", Journal of Basic and Clinical Physiology and Pharmacology, 2020

Publication

<1 %

11

Aji Prasetyaningrum, Bangkit Suryo Wicaksono, Azafilmi Hakiim, Aulia Dwi Ashianti et al. "Ultrasound-Assisted Encapsulation of Citronella Oil in Alginate/Carrageenan Beads: Characterization and Kinetic Models", ChemEngineering, 2023

Publication

<1 %

12

Hayder A. Hammoodi. "Moringa and its importance in human life", Science Archives, 2021

Publication

<1 %

13

Hefeng Wang, Anran Xu, Liping Gong, Zhaowen Chen, Bin Zhang, Xiuyun Li. "The Microbiome, an Important Factor That Is Easily Overlooked in Male Infertility", Frontiers in Microbiology, 2022

Publication

<1 %

14

Asmaa Abdelsalam, Kamal Chowdhury, Arezue Boroujerdi, Ahmed El-Bakry. "Effect of stress hormones on the metabolome of a suspension culture of the aromatic medicinal plant *Cymbopogon schoenanthus* subsp. *proximus*", Plant Cell, Tissue and Organ Culture (PCTOC), 2023

Publication

<1 %

15

Alessandra Gallo, Maria Consiglia Esposito, Raffaele Boni, Elisabetta Tosti. "Oocyte quality assessment in marine invertebrates: a novel approach by fluorescence spectroscopy", Biological Research, 2022

Publication

<1 %

16

V Hasan, A Wijayanti, M B Tamam, R A Islamy, M S Widodo. "Beardless barb *Cyclocheilichthys apogon* (Valenciennes, 1842) (Cypriniformes, Cyprinidae):

<1 %

Distribution extension and first record from South Bali", IOP Conference Series: Earth and Environmental Science, 2021

Publication

17

Zubeir Ali, James Broome, Ian Pearce. "The History and clinical examination for self-reported erectile dysfunction: are we following European guidelines? The results of the North West of England Regional Audit", Journal of Clinical Urology, 2016

Publication

<1 %

18

Deenadayalan Karaiyagowder Govindarajan, Vijayakumar Selvaraj, Anit Selva Jobitha Mariya Selvaraj, Sabeena Shahul Hameed et al. "Green synthesis of silver micro- and nanoparticles using phytochemical extracts of Cymbopogon citratus exhibits antibacterial properties", Materials Today: Proceedings, 2022

Publication

<1 %

19

Fatemeh Ghasemian, Shahin Esmaeilnezhad, MohammadJavad MehdipourMoghaddam. "Staphylococcus saprophyticus and Escherichia coli: Tracking from sperm fertility potential to assisted reproductive outcomes", Clinical and Experimental Reproductive Medicine, 2021

Publication

<1 %

20

Samy Hussein, Samir Abdel-Aal, Shalabia Emam. "Phytochemical analysis and Antiproliferative activity of lotus peregrinus L. against MCF7 (breast carcinoma cell line)", Benha Veterinary Medical Journal, 2019

Publication

<1 %

21

Yong Zhang, Xian-Li Song, Bin Yu, Lian-Chee Foong, Yu Shu, Chun-Wai Mai, Jing Hu, Baijun Dong, Wei Xue, Chee Wai Chua. " loss-of-function causes vulnerability to autophagy inhibition in aggressive prostate cancer ", International Journal of Urology, 2022

Publication

<1 %

22

Huifeng Luo, Rongxiu Liu, Yilin Lang, Jinhui Zhao, Cuicui Zhuang, Jundong Wang, Chen Liang, Jianhai Zhang. "Melatonin alleviated fluoride-induced impairment of spermatogenesis and sperm maturation process via Interleukin-17A", Food and Chemical Toxicology, 2023

Publication

<1 %

23

A Zikopoulos, S Grigoriadis, D Karagkouni, E Maziotis et al. "O-137 Mapping microRNA profiling and dysregulated pathways in non-obstructive azoospermia cases with maturation arrest: A systematic review and in-silico analysis of the affected pathways", Human Reproduction, 2023

<1 %

24

Ying Chen, Cheng Cheng, Wenqing Xu, Yanfan Cui et al. "The risk of micro/nanoplastics to human sperm and the intervention against their toxicity using magnetic Fe₃O₄ nanoparticles", Research Square Platform LLC, 2023

Publication

<1 %

25

Mallol, Anna, Josep Santaló, and Elena Ibáñez. "Improved Development of Somatic Cell Cloned Mouse Embryos by Vitamin C and Latrunculin A", PLoS ONE, 2015.

Publication

<1 %

26

Hao Wang, Jiwei Zhang, Dongyue Ma, Ziwei Zhao, Bin Yan, Fu Wang. "The role of red ginseng in men's reproductive health: a literature review", Basic and Clinical Andrology, 2023

Publication

<1 %

27

Sonia Gupta, Vijay Prabha. " Human Sperm Interaction with : A Molecular Approach ", Journal of Pathogens, 2012

Publication

<1 %

28

Sulagna Dutta, Pallav Sengupta, Emmanuel Izuka, Ifeanyi Menuba, Ravindran Jegasothy, Uchenna Nwagha. "Staphylococcal infections and infertility: mechanisms and

<1 %

management", Molecular and Cellular Biochemistry, 2020

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off

FINAL GRADE

GENERAL COMMENTS

/0

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU *PEER REVIEW*
KARYA ILMIAH : JURNAL ILMIAH

Judul Jurnal Ilmiah (Artikel) : *"Examining the Effects of Lemongrass (Cymbopogon nardus) Extract on Sperm Quality and Testosterone Level in Mice Induced with Staphylococcus aureus"*

Jumlah Penulis : 6 orang

Status Pengusul : ~~Penulis Utama~~/Penulis ke-2 /Penulis korespondensi

Identitas Jurnal Ilmiah :

- a. Nama Jurnal : RJPT: Research Journal of Pharmaceutical and Technology
- b. Nomor ISSN : Online ISSN: 0974-360X
Print ISSN: 0974-3618
- c. Nomor/Volume : Volume 17 / Nomor 02
- d. Edisi (bulan/tahun) : Februari, 2024
- e. Penerbit : A and V Publication
- f. DOI artikel :
<https://www.rjptonline.org/AbstractView.aspx?PID=2024-17-2-64>
- g. Alamat Web Jurnal :
<https://www.rjptonline.org/AboutJournal.aspx>
- h. Jumlah halaman : 6 Halaman

: ☒ Jurnal Internasional Terindeks basis data internasional bereputasi dan berimpact faktor

☐ Jurnal Nasional Terakreditasi/ Peringkat 1 dan 2 SINTA

☐ Jurnal Nasional DOAJ/CABI/COPERNICUS/Peringkat 3 dan 4 SINTA

☐ Jurnal Nasional Peringkat 5 dan 6 SINTA

Hasil Penilaian *Peer Review* :

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah				Nilai Akhir Yang Diperoleh (NA)
	Jurnal Internasional terindeks basis data internasional bereputasi <input checked="" type="checkbox"/>	Jurnal Nasional Terakreditasi. Peringkat 1 dan 2 SINTA <input type="checkbox"/>	Jurnal Nasional DOAJ/CABI/Copernicus/Peringkat 3 dan 4 SINTA <input type="checkbox"/>	Jurnal Nasional Peringkat 5 dan 6 SINTA <input type="checkbox"/>	
e. Kelengkapan unsur isi artikel (10%)	4	2,5	2	1,5	3,6
f. Ruang lingkup dan kedalaman pembahasan (30%)	12	7,5	6	4,5	11,7
g. Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)	12	7,5	6	4,5	12
h. Kelengkapan unsur dan kualitas terbitan/jurnal (30%)	12	7,5	6	4,5	12
Total = (100%)	40	25	20	15	39,3
Nilai Pengusul (NA x BP)	39,3 x 0,4=		15,72		

Catatan Penilaian Kualitatif oleh Reviewer:

Kelengkapan unsur dan isi artikel terpenuhi, ruang lingkup dan kedalaman pembahasan sudah baik, metodologi yang digunakan cukup mutakhir, kelengkapan dan kualitas jurnal baik dan sudah terakreditasi internasional bereputasi. Artikel masih berstatus diterima dan terpublish pada bulan februari. Kesesuaian isi artikel dengan bidang ilmu pengusul sudah sesuai.

Catatan Bobot Pengusul:

Sendiri = 1; Penulis pertama + korespondensi = 0,6;
 Penulis pertama = 0,4; Penulis korespondensi = 0,4
 Anggota = 0,4 atau 0,2 dibagi banyak anggota

Lamongan, 10 Januari 2024

Reviewer 1,



Prof. Win Darmanto, Ph.D

NIP. 196106161987011001

Unit kerja : Dep. Biologi, FST, UNAIR

Jabatan Akademik Terakhir: Guru Besar

Bidang Ilmu : Biologi/ Fisiologi Hewan

LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU *PEER REVIEW*
KARYA ILMIAH : JURNAL ILMIAH

Judul Jurnal Ilmiah (Artikel) : *"Examining the Effects of Lemongrass (Cymbopogon nardus) Extract on Sperm Quality and Testosterone Level in Mice Induced with Staphylococcus aureus"*

Jumlah Penulis : 6 orang

Status Pengusul : Penulis Utama/ Penulis ke-2 / Penulis korespondensi

Identitas Jurnal Ilmiah :

- a. Nama Jurnal : RJPT: Research Journal of Pharmaceutical and Technology
- b. Nomor ISSN : Online ISSN: 0974-360X
Print ISSN: 0974-3618
- c. Nomor/Volume : Volume 17 / Nomor 02
- d. Edisi (bulan/tahun) : Februari, 2024
- e. Penerbit : A and V Publication
- f. DOI artikel :
<https://www.rjptonline.org/AbstractView.aspx?PID=2024-17-2-64>
- g. Alamat Web Jurnal :
<https://www.rjptonline.org/AboutJournal.aspx>
- h. Jumlah halaman : 6 Halaman

☒ Jurnal Internasional Terindeks basis data internasional bereputasi dan berimpact faktor
☐ Jurnal Nasional Terakreditasi/ Peringkat 1 dan 2 SINTA
☐ Jurnal Nasional DOAJ/CABI/COPERNICUS/Peringkat 3 dan 4 SINTA
☐ Jurnal Nasional Peringkat 5 dan 6 SINTA

Hasil Penilaian *Peer Review* :

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah				Nilai Akhir Yang Diperoleh (NA)
	Jurnal Internasional terindeks basis data internasional bereputasi <input checked="" type="checkbox"/>	Jurnal Nasional Terakreditasi. Peringkat 1 dan 2 SINTA <input type="checkbox"/>	Jurnal Nasional DOAJ/CABI/Co pernicus/Peringkat at 3 dan 4 SINTA <input type="checkbox"/>	Jurnal Nasional Peringkat 5 dan 6 SINTA <input type="checkbox"/>	
a. Kelengkapan unsur isi artikel (10%)	4	2,5	2	1,5	3,7
b. Ruang lingkup dan kedalaman pembahasan (30%)	12	7,5	6	4,5	12
c. Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)	12	7,5	6	4,5	12
d. Kelengkapan unsur dan kualitas terbitan/jurnal (30%)	12	7,5	6	4,5	11,7
Total = (100%)	40	25	20	15	39,4
Nilai Pengusul (NA x BP)	39,4 x 0,4=		15,76		

Catatan Penilaian Kualitatif oleh Reviewer:

Kelengkapan unsur isi artikel sudah sesuai, ruang lingkup dan kedalaman pembahasan baik dan menggunakan literatur yang terkini sesuai dengan topik penelitian. Metode yang digunakan menggunakan sumber yang sesuai. Artikel sudah diterima pada jurnal internasional bereputasi terindeks scopus Q2.

Catatan Bobot Pengusul:

Sendiri = 1; Penulis pertama + koresponding = 0,6;
 Penulis pertama = 0,4; Penulis koresponding = 0,4
 Anggota = 0,4 atau 0,2 dibagi banyak anggota

Lamongan, 08 Januari 2024

Reviewer 2,



Dr. Nastiti Intan Permata Sari, S.Si., M.Ked.Trop

NIDN. 4720069301

Unit kerja: Biologi, FMIPA Militer, UNHAN

Jabatan Akademik Terakhir: Lektor

Bidang Ilmu: Biologi/ Biologi Molekuler



A AND V PUBLICATIONS

Article's Basic Information:

Paper ID : 221128224141309810

Paper Title : Examining the Effects of Lemongrass (*Cymbopogon nardus*) Extract on Sperm Quality and Testosterone Level in Mice Induced with *Staphylococcus aureus*

Authors : Lilis Maghfuroh, Putri Ayu Ika Setiyowati, Rofiatun Solekha, Nynda Ayu Nadira Savitri, Eka Febrianti Wulandari

Author's Email : lilisahza99@gmail.com; putriayuikasetiyowati@gmail.com; rofiatunsolekha22@gmail.com; angellaananda466@gmail.com

Submitted to Journal : Research Journal of Pharmacy and Technology

Submitted By : putri ayu ika setiyowati (putriayuikasetiyowati@gmail.com)

Date of Submission : 28 November, 2022

Comments From Reviewer:



Viol Dhea Kharisma

To begin with, this manuscript is acceptable based on the Turnitin plagiarism check. In addition, in order to improve the readability of the paper by an English editing service. Furthermore, the paper is easy to follow by the reader. However, I may require some comments on the following issues.

Title: The title is good.

Abstract: This section was well-written and easy to understand.

Introduction: Adequate.

Material and Methods: Okay

Results: Good

Discussion: The discussion of the study lacks information about the important topics.

Conclusion: Good.

References: Updated references are needed.



Arif Nur Muhammad Ansori

- Is the work clearly and accurately presented and does it cite the current literature? Yes
 - Is the study design appropriate and is the work technically sound? Yes
 - Are the conclusions drawn adequately supported by the results? Yes
- Overall, the manuscript is good and easy to follow. But, they need improvement for English editing (native English services).

Comments From Editorial Board:



Resubmission of Article

Dear Author,
 Pl Resubmit the article after making corrections suggested by Reviewers.
 Thanks
 Editor

**Editorial Board Team**

In order to improve the quality of the paper you just go through the below link and cite atleast 10 articles from the list, that have similar works as of yours. The link is <http://www.anvpublication.org/ViewReviewComments.aspx?tmppid=221128224141309810> .

[Print Report](#)



A and V Publications

Report Generated on: 1/25/2024 6:33:22 AM

BASIC INFORMATION

Paper ID: **221128224141309810**

Submission Date: **November 28, 2022**

Paper Title: **Examining the Effects of Lemongrass (*Cymbopogon nardus*) Extract on Sperm Quality and Testosterone Level in Mice Induced with *Staphylococcus aureus***

Author(s) Name: Lilis Maghfuroh, Putri Ayu Ika Setiyowati, Rofiatun Solekha, Nynda Ayu Nadira Savitri, Eka Febrianti Wulandari

Author(s) Email: lilisahza99@gmail.com; putriayuikasetiyowati@gmail.com; rofiatunsolekha22@gmail.com; angellaananda466@gmail.com

Author(s) Address: Nursing Program, Faculty of Health Science, Universitas Muhammadiyah Lamongan, East Java, Indonesia, Biology Program, Faculty of Science, Technology, and Education, Universitas Muhammadiyah Lamongan, East Java, Indonesia, Biology Program, Faculty of Science, Technology, and Education, Universitas Muhammadiyah Lamongan, East Java, Indonesia, Biology Program, Faculty of Science, Technology, and Education, Universitas Muhammadiyah Lamongan, East Java, Indonesia, Nursing Program, Faculty of Health Science, Universitas Muhammadiyah Lamongan, East Java, Indonesia,

Journal: Research Journal of Pharmacy and Technology

Submitted By: putri ayu ika setiyowati

Email ID: putriayuikasetiyowati@gmail.com

REVIEWER INFORMATION

	First Reviewer	Second Reviewer
Name:	Arif Nur Muhammad Anshori	Viol Dhea Kharisma
Email ID:	ansori.anm@gmail.com	violdhea@gmail.com
Mobile No:	082144647832	08121787002
Address:	Universitas Airlangga	Universitas Airlangga

PAYMENT DETAILS

Order No.:	202343114323696510870008	Order Date:	April 03, 2023, 11:04
Amount INR:	3540.00	OR Amount USD:	210.00
Payment Date:	April 03, 2023	Transaction No.:	pay_LZDSfAuiTirJ7K
Payment Status:	PAYMENT VERIFIED		

PAPER PUBLICATION / PROCESSING STATUS

- ▶▶ 28/Nov/2022, 10:41:41 PM --- Article submitted by the author.
- ▶▶ 30/Nov/2022, 09:39:14 AM --- Article sent back to author for minor corrections.
- ▶▶ 30/Nov/2022, 09:39:14 AM --- New comments from editorial board.
- ▶▶ 08/Dec/2022, 11:23:37 PM --- Article resubmitted by author after correction.
- ▶▶ 11/Jan/2023, 06:38:51 PM --- Article is sent to reviewers.
- ▶▶ 09/Feb/2023, 05:17:48 AM --- Review comments submitted by the reviewer.
- ▶▶ 09/Feb/2023, 05:18:09 AM --- Review comments submitted by the reviewer.
- ▶▶ 10/Feb/2023, 12:49:21 PM --- Article sent back to author for minor corrections.
- ▶▶ 10/Feb/2023, 12:49:28 PM --- New comments from editorial board.
- ▶▶ 18/Feb/2023, 07:17:24 PM --- Article resubmitted by author after correction.
- ▶▶ 31/Mar/2023, 06:48:20 PM --- Final version of article is required.
- ▶▶ 03/Apr/2023, 05:17:23 AM --- Final version of article submitted by author.
- ▶▶ 03/Apr/2023, 05:17:23 AM --- Article is accepted by publisher.
- ▶▶ 03/Apr/2023, 11:43:23 AM --- Payment proof required.
- ▶▶ 03/Apr/2023, 01:00:24 PM --- Payment proof uploaded by author.
- ▶▶ 06/Apr/2023, 03:14:16 PM --- Payment details verified by publisher.
- ▶▶ 08/Apr/2023, 04:39:37 PM --- Payment details verified by publisher.