

Determinants of Iron Supplementation (Fe) Adherence among Pregnant Women at Puskesmas Sugio Lamongan

by Te Mason

Submission date: 29-Feb-2024 10:34PM (UTC-0500)

Submission ID: 2306933498

File name: I_Diah_Eko_Des_2021_The_Indonesian_Journal_of_Health_Science.pdf (242.66K)

Word count: 3573

Character count: 19486

Determinants of Iron Supplementation (Fe) Adherence among Pregnant Women at Puskesmas Sugio Lamongan

Diah Eko Martini^{*1}, Heny Ekawati¹, Lilis Maghfuroh¹, Silvy Harmiadillah¹,
Wahyu Retno Gumelar¹, Aizatun Nisa¹

¹Department of Maternity Nursing Program, Faculty of Health Science, Universitas Muhammadiyah Lamongan

*Corresponding author: Diah Eko Martini, Email: diahekomartini@gmail.com

Received: 21 November 2021 | Accepted: 21 December 2021 | Published: 30 December 2021

Abstract

Background and Aim: Anemia during gestation is a complication that often occurs in pregnant women. This condition can be controlled by providing iron (Fe) supplementation; therefore, the government has set a policy of minimum service standards for the coverage of 90 tablets of iron (Fe) supplementation for the period of pregnancy. However, pregnant women's adherence to iron (Fe) supplementation is still low. The objective of this research was to determine the factors that influence pregnant women's adherence to iron (Fe) supplementation.

Methods: The design of this study used a cross-sectional study approach. The sample was used for all pregnant women who were recorded at the Sugio Lamongan Health Center in April 2021 totaling 36 respondents, the data was taken using a structured questionnaire and analyzed using chi-square and Fisher's Exact Test.

Results: The findings revealed that the factors that influence pregnant women's adherence to iron (Fe) supplementation are knowledge, family support, regularity of the ANC. Pregnant women (66.6%) were obedient in taking iron (Fe) supplements (PR=0.227) is a factor of pregnant women's adherence to taking iron (Fe) supplements.

Conclusion: Therefore, all access that affects antenatal care services and follow-up ANC services is an important element to improve the Adherence of pregnant women in taking iron (Fe) supplementation.

Keywords: Anemia; Antenatal care; Family; Iron; Knowledge; Pregnant Women

How to cite this article: Martini, et. al. (2021). Determinants of iron supplementation (Fe) adherence among pregnant women at Puskesmas Sugio Lamongan. *The Indonesian Journal of Health Science*, 13(2), 180-188. DOI: 10.32528/ijhs.v13i2.6426.

Copyright: ©2021 Martini, et. al. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Published by: Universitas Muhammadiyah Jember
ISSN (Print): 2087-5053
ISSN (Online): 2476-9614

INTRODUCTION

Anemia during pregnancy is a condition where there is a decrease in hemoglobin or the number of red blood cells less than 11g% in the I trimester and III trimester or levels of less than 10.5g% in the II trimester (Novitasari et al., 2019). Pregnancy anemia occurs due to an increase in body fluids (plasma fluids) that are not comparable to the addition of blood cells so that blood dilution (Hemodilution) occurs. Pregnant women are susceptible to anemia. Iron insufficiency causes anemia in 41.8 percent of pregnant women around the world (Ministry of Health, 2014).

The prevalence of pregnant women with anemia in Indonesia was 37.1% while in 2018 the number increased by 48.9%. This number exceeds severe public health problems whose prevalence limit as 40%. The average coverage of national iron supplementation is 85.1% and the average coverage of iron supplementation in East Java Province in 2018 is still below the average of 90.82% with a target 90% this means there is still a gap of 2.9%. Based on data from the Nutrition Section of the Lamongan District Office in 2018, the average coverage of iron supplement in Lamongan District is still low. While the results of the preliminary study interview with 21 pregnant women for 2 days on February 04-05, 2021 who checked their pregnancy and who live around Sugio Health Center, obtained the results that only 9 (45%) of pregnant women obey to consume iron supplementation (Fe), while 12 (55%) of pregnant women do not comply to consume iron supplementation. Anemia may result from several causes, with the most significant contributor being iron deficiency (World Health Organization, 2015).

Iron deficiency anemia can cause adverse perinatal outcomes, such as miscarriage, bleeding, stillbirth, fetal growth retardation, low birth weight, premature birth, and neonatal death (Beckert et al., 2019; Hare et al., 2013; Tunkyi & Moodley, 2018). Similarly, Begum, (2012) and Ibrahim et al., (2013) stated that anemia for the period of pregnancy has fetal and neonatal risks, which include prematurity, low birth weight, and fetal distress, which contribute to illness and perinatal mortality.

Pregnant woman needs a large iron (1000 mg) during pregnancy is not enough if obtained from food only, therefore, it should be helped by iron (Fe) supplementation (Ministry of Health, 2014). Therefore, the central government set a minimum service standard of iron supplementation coverage (Fe) during pregnancy as much as 90 tablets for the prevention of anemia in pregnancy. Iron supplementation is an effective way due to the content of iron is equipped with folic acid that prevent anemia, but the adherence of pregnant women to consume iron supplements is still low. Non-adherence of pregnant women consuming iron supplements (Fe) can have a great chance of developing anemia (Kautshar & Jafar, 2013). Several factors could influence pregnant women's willingness to take iron supplements (Fe) including knowledge, motivation, family support, and antenatal care visit (Kamidah, 2015). The condition of non-adherence in consuming iron tablets is one of the causes of anemia in pregnant women (Hidayah & Anasari, 2012). Pregnant women are one of the groups at high risk for nutritional problems, especially iron deficiency anemia, because the pregnant women need for iron (Fe)

increases significantly during pregnancy. One of the factors causing the high iron deficiency anemia in pregnant women is the low adherence of pregnant women in consuming Fe tablets (Kadir, 2019). Iron tablets are mineral tablets needed by the body for the formation of red blood cells or hemoglobin. If humans lack iron in the food they eat every day, it can cause nutritional anemia (lack of blood). Iron (Fe) tablets are needed by pregnant women, so pregnant women are required to take at least 60 tablets of Fe during pregnancy. Iron (Fe) acts as a component that forms myoglobin, a protein that distributes oxygen to muscles, forms enzymes, and collagen. In addition, iron also plays a role in the body's resistance. Iron (Fe) tablets are important for pregnant women because they have several functions, including: Increasing nutritional intake to the fetus, Preventing iron deficiency anemia, Preventing bleeding during childbirth, Reducing the risk of death in the mother due to bleeding during childbirth (Nugraheni, 2020).

This condition of iron deficiency will increase the risk of anemia which can cause bleeding, attached placenta, uterine tears that loosen due to frequent childbirth (Abu-Ouf & Jan, 2015). Therefore, it is necessary to increase the compliance of pregnant women in consuming iron (Fe) supplements, this can be done in various ways, among others by conducting counseling about the importance of consuming iron (Fe) supplements, optimizing Maternal and Child Health Services in every community service such as posyandu and puskesmas in each region, surveying mothers' knowledge about the importance of consuming iron (Fe) supplements, intervening in pregnant women, improving the skills of health workers in each area, and empowering

others or community shops to know the importance of consuming iron supplements. iron (Fe) during pregnancy. In addition, effective interaction or communication between health workers and pregnant women is very important. This determines success in solving health problems. This effective interaction can reduce patient doubts, and increase patient compliance in taking supplements iron (Fe) (Endang Fourianalistyawati, 2012).

METHODS

The research design was used a cross-sectional study with the population being pregnant women who had a pregnancy check-up at the Sugio Health Center, Sugio District, Lamongan Regency during April 2021. Furthermore, the sampling in this study was used a consecutive sampling technique with 36 respondents. Data retrieval using questionnaires that have been tested for validity and reliability then analyzed with *chi-square* test. The data was analyzed using the program statistical package of socio science (SPSS) version 24.0 with a value of 0.05.

This research has been reviewed and received approval from the Health research ethics commission of Muhammadiyah Lamongan University with Number 121 / EC /KEPK - S2 / 01 / 2021.

RESULTS

Table 1 shows that the majority of pregnant women aged 20-35 years (69.4%), while based on education most pregnant women have a high school education background (91.7%). While based on gestational age the majority of pregnant women are still entering pregnancy Trimester II (63.9%).

Table 1. Demographic Characteristics of Pregnant Women in Sugio Lamongan Health Center

Characteristic	Frequency	Percentage %
Age (year)		
<20	6	16.6
20-35	25	69.4
>35	5	13.8
Total	36	100
Education		
Junior high school	0	0
Senior high school	33	91.7
Bachelor	3	8.3
Total	36	100
Occupation		
Factory workers	8	22.2
Teacher	2	5.5
Housewife	15	41.7
Self-employed	11	30.6
Total	36	100
Gestational Age		
0-12 weeks	3	8.3
13-28 weeks	23	63.9
29-40 weeks	10	27.8
Total	36	100

Table 2. Determining Factors of Adherence of Pregnant Women in taking Iron (Fe) Supplements

Characteristic	N	%	PR	95%CI	p-value
Adherence					
Obedient	24	66.6			
Disobedient	12	33.3			
Knowledge					
Good	26	72.3	4.857	1.096 - 21.517	0.031
Less	10	27.7			
Motivation					
Good	33	91.7	1.222	0.863 - 1.731	0.307
Less	3	8.3			
Family support					
Good	17	47.2	0.292	0.156 - 0.544	0.000
Less	19	52.8			
Regularity of ANC					
Regular	20	55.6	0.227	0.102 - 0.505	0.000
Irregular	16	44.4			

Table 2 shows that the majority (66.6%) of pregnant women comply with consuming iron supplements (Fe). Pregnant women with good knowledge have 4.85 times more possible to comply with iron supplements (Fe) for

90 days (PR=4,857; [95%CI:1.096-21.517]), while pregnant women who have good family support are 0.292 times more obedient in taking iron supplements (Fe) for 90 days (PR=0.292; [95%CI:0.156-0.544]). Similarly, pregnant women who regularly visited the ANC had a 0.227 chance of complying with iron supplement consumption (Fe) for 90 days compared to irregular pregnant women in ANC visits (OR=0.227; [95%CI: 0.102-0.505]). Based on the results above, it can be concluded that the factors of knowledge, family support, and regularity in an ANC visit are the determining factors of adherence of pregnant women in taking iron supplements (Fe) for 90 days. As for the motivation, a factor is known not to be a factor that affects Adherence in iron supplements (Fe) for 90 days (P-value = 0.307; [95%CI: 0.863 - 1.731]).

DISCUSSION

Iron is essential in pregnancy and infancy to chance the need for hematopoiesis, growth, and development (Brannon & Taylor, 2017). Adherence of pregnant women in taking iron supplementation is important in the prevention of pregnancy anemia. Adherence with iron supplements (Fe) refers to a person's consistency in taking iron supplements (Fe) as prescribed by a doctor, which is a daily dose of 1 tablet (60 mg elemental iron and 0.25 g folic acid) for at least 90 days during pregnancy (Ministry of Health, 2014).

Pregnant women who follow the recommendation of health professionals to use Fe Tablets are said to be adhering to the advice. Because iron tablets have a solid iron content and are combined with folic acid, they are regarded an efficient method of

supplementing (Parulian, 2018). The results found that the majority of pregnant women adherence to the consumption of iron supplements (Fe). Crosstab results showed that mothers who have good years and have good motivation tend to be obedient in taking iron supplements (Fe), as well as pregnant women who have good family support and regularly in an ANC visit also tend to be obedient in consuming iron supplements (Fe). These results were reinforced by the *chi-square* test and Fisher's Exact Test which stated that the variable knowledge, family support, and regularity of ANC visits are the determining factors and have a significant opportunity to assess whether or not pregnant women take iron supplementation (Fe).

Knowledge of iron supplementation (Fe) and its benefits become one of the factors that encourage mothers to comply in consuming iron supplements (Fe) and the majority of pregnant women who consume iron supplements (Fe) know the benefits and purposes of consuming iron (Fe) (Wiradnyani et al., 2013). A good pregnant woman's knowledge of the importance of consuming iron supplements (Fe) will allow pregnant women to analyze and synthesize information obtained to improve their well-being and family through positive behaviors, in which case pregnant women will try to avoid the risk of anemia during pregnancy because she can predict risks that may harm herself and her fetus. Conversely, if the knowledge is low, then it is likely to refuse to take iron supplements (Fe) regularly, especially if it feels that there are annoying side effects. Knowledge is closely related to the level of education where most pregnant women are educated in high

school. A study conducted by (Ali et al., 2021) stated that educated and well-educated women are usually more concerned about the health of their future babies compared to uneducated women.

Family support is also a determining factor in adherence to consuming iron supplements (Fe). Triharini et al. (2018) stated that family support is one of the factors that affect pregnant women's Adherence to taking iron supplements (Fe). Family in advising on the problem of taking iron tablets can also be advice and encouragement so as not to consider discomfort due to iron supplementation as a hindrance. Fe tablet adverse effects, such as nausea, can make it difficult for pregnant women to take iron supplements regularly Ali et al., (2021) . The family can provide adherence support for iron supplementation by directly monitoring its pregnant relatives daily.

Regular ANC visits are also noted as a correlated factor to the adherence of pregnant women taking iron supplements (Fe). Another study from Lacerte et al., (2011) also stated that the number of prenatal visits confirmed by women in consuming iron (Fe) supplements, even Getachew et al., (2018) stated that the number of ANC visits of four times or more significantly affected adherence. Antenatal care is an important access for compliant pregnant women to consume iron supplements (Fe), this is possible because health care providers can help pregnant women during visitation so that they can discuss everything related to pregnancy anemia and its relationship with iron supplements (Fe) and encourage them to take tablets according to prescription and provide health education about the benefits of taking iron supplements

(Fe) during pregnancy. and other study results also state that knowledge, family support, and the number of prenatal visits are factors that correlate to the adherence of women who pregnant to taking iron supplementation (Fe). Anggraini et al., (2018) stated that the better-quality interaction of women who pregnant with health workers, the more likely pregnant women are to obediently take iron supplements (Fe) to prevent anemia during pregnancy.

CONCLUSION

The adherence of pregnant women to consuming iron supplements (Fe) is still low. Factors of knowledge, family support, and regularity of the ANC are factors that are recorded significantly in determining the adherence of pregnant women to taking iron supplements (Fe) for the period of pregnancy.

REFERENCES

- Abu-Ouf, N. M., & Jan, M. M. (2015). The impact of maternal iron deficiency and iron deficiency anemia on child's health. *Saudi Medical Journal*, 36(2), 146–149.
<https://doi.org/10.15537/smj.2015.2.10289>.
- Ali, S. A., Ali, S. A., Razzaq, S., Khowaja, N., Gutkind, S., Raheman, F. U., & Suhail, N. (2021). Predictors of iron consumption for at least 90 days during pregnancy: Findings from National Demographic Health Survey, Pakistan (2017-2018). *BMC Pregnancy and Childbirth*, 21(1), 352.
<https://doi.org/10.1186/s12884-021-03825-2>.
- Anggraini, D. D., Purnomo, W., & Trijanto, B. (2018). *Interaksi*

- Ibu Hamil Dengan Tenaga Kesehatan Dan Pengaruhnya Terhadap Kepatuhan Ibu Hamil Mengonsumsi Tablet Besi (Fe) Dan Anemia Di Puskesmas Kota Wilayah Selatan Kota Kediri. *Buletin Penelitian Sistem Kesehatan*, 21(2), 92–89.
<https://doi.org/10.22435/hsr.v2i2.346>.
- Beckert, R. H., Baer, R. J., Anderson, J. G., Jelliffe-Pawlowski, L. L., & Rogers, E. E. (2019). Maternal anemia and pregnancy outcomes: A population-based study. *Journal of Perinatology: Official Journal of the California Perinatal Association*, 39(7), 911–919.
<https://doi.org/10.1038/s41372-019-0375-0>.
- Begum, S. (2012). Factors associated with adherence to iron folic acid supplementations during pregnancy in Uttar Pradesh. Undefined.
<https://www.semanticscholar.org/paper/Factors-associated-with-adherence-to-iron-folic-in-Begum/aedcdda59968a806877dd5206679b4d7127993f6>.
- Brannon, P. M., & Taylor, C. L. (2017). Iron Supplementation during Pregnancy and Infancy: Uncertainties and Implications for Research and Policy. *Nutrients*, 9(12), E1327.
<https://doi.org/10.3390/nu9121327>.
- Endang Fourianalisyawati, M. P. (2012). Komunikasi Yang Relevan Dan Efektif Antara Dokter Dan Pasien. *Jurnal Psikogenesis*, 1(1), 82–87.
<https://doi.org/10.24854/jps.v1i1.37>.
- Getachew, M., Abay, M., Zelalem, H., Gebremedhin, T., Grum, T., & Bayray, A. (2018). Magnitude and factors associated with adherence to Iron-folic acid supplementation among pregnant women in Eritrean refugee camps, northern Ethiopia. *BMC Pregnancy and Childbirth*, 18(1), 83.
<https://doi.org/10.1186/s12884-018-1716-2>.
- Hare, G. M. T., Freedman, J., & David Mazer, C. (2013). Review article: Risks of anemia and related management strategies: can perioperative blood management improve patient safety? *Canadian Journal of Anaesthesia = Journal Canadien D'anesthésie*, 60(2), 168–175.
<https://doi.org/10.1007/s12630-012-9861-y>.
- Hidayah, W., & Anasari, T. (2012). Hubungan Kepatuhan Ibu Hamil Mengonsumsi Tablet Fe Dengan Kejadian Anemia Di Desa Pageraji Kecamatan Cilongok Kabupaten Banyumas. *Bidan Prada : Jurnal Publikasi Kebidanan STIKes YLPP Purwokerto*, 3(02).
<https://ojs.stikesylpp.ac.id/index.php/Prada/article/view/59>.
- Ibrahim, Z. M., El-Hamid, S. A., Mikhail, H., & Khattab, M. (2013). Assessment of Adherence to Iron and Folic Acid Supplementation and Prevalence of Anemia in Pregnant Women.
<https://www.semanticscholar.org/paper/Assessment-of-Adherence-to-Iron-and-Folic-Acid-and-Ibrahim-El-Hamid/9477a6627d131b008e2dbfb71fa4fcd146d51074>.

- Kadir, S. (2019). Faktor Penyebab Anemia Defisiensi Besi Pada Ibu Hamil Di Wilayah Kerja Puskesmas Bongo Nol Kabupaten Boalemo. *Jambura Journal of Health Sciences and Research*, 1(2), 54–63. <https://doi.org/10.35971/jjhsr.v1i2.2396>.
- Kamidah. (2015). Faktor- Faktor Yang Mempengaruhi Kepatuhan Ibu Hamil Mengonsumsi Tablet Fe Di Puskesmas Simo Boyolali. *Gaster*, 12(1), 36–45.
- Kautshar, N., & Jafar, N. (2013). Kepatuhan Ibu Hamil Dalam Mengonsumsi Tablet Zat Besi (Fe) Di Puskesmas Bara-Baraya Tahun 2013. 15.
- Lacerte, P., Pradipasen, M., Temcharoen, P., Imamee, N., & Vorapongsathorn, T. (2011). Determinants of Adherence to Iron/Folate Supplementation During Pregnancy in Two Provinces in Cambodia. *Asia Pacific Journal of Public Health*, 23(3), 315–323.
- Ministry of Health. (2014). Pusat Data dan Informasi—Kementerian Kesehatan Republik Indonesia. Profil Kesehatan Indonesia Tahun 2014. <https://pusdatin.kemkes.go.id/article/view/15060500001/profil-kesehatan-indonesia-tahun-2014.html>.
- Novitasari, Y. D., Wahyudi, F., & Nugraheni, A. (2019). Faktor – Faktor Yang Berhubungan Dengan Kekurangan Energi Kronik (Kek) Ibu Hamil Di Wilayah Kerja Puskesmas Rowosari Semarang. *Diponegoro Medical Journal (Jurnal Kedokteran Diponegoro)*, 8(1), 562–571. <https://doi.org/10.14710/dmj.v8i1.23399>.
- Nugraheni, A. S. (2020). Literature Review: Pengaruh Penyuluhan Terhadap Pengetahuan Ibu Hamil Trimester I Tentang Tablet FE [Diploma, UNIMUS]. <http://repository.unimus.ac.id/4653/>.
- Parulian, I. (2018). Strategi Dalam Penanggulangan Pencegahan Anemia Pada Kehamilan. *Jurnal Ilmiah Widya*, 4(3), Article 3. <https://ejournal.jurwidyakop3.com/index.php/jurnal-ilmiah/article/view/255>.
- Triharini, M., Nursalam, Sulistyono, A., Adriani, M., Armini, N. K. A., & Nastiti, A. A. (2018). Adherence to iron supplementation amongst pregnant mothers in Surabaya, Indonesia: Perceived benefits, barriers and family support. *International Journal of Nursing Sciences*, 5(3), 243–248. <https://doi.org/10.1016/j.ijnss.2018.07.002>.
- Tunky, K., & Moodley, J. (2018). Anemia and pregnancy outcomes: A longitudinal study. *The Journal of Maternal-Fetal & Neonatal Medicine: The Official Journal of the European Association of Perinatal Medicine, the Federation of Asia and Oceania Perinatal Societies, the International Society of Perinatal Obstetricians*, 31(19), 2594–2598. <https://doi.org/10.1080/14767058.2017.1349746>.
- Wiradnyani, L. A. A., Khusun, H., & Achadi, E. L. (2013). Faktor-

Faktor Yang Berhubungan Dengan Kepatuhan Ibu Mengonsumsi Tablet Besi-Folat Selama Kehamilan. *Jurnal Gizi Dan Pangan*, 8(1), 63–70.
<https://doi.org/10.25182/jgp.2013.8.1.63-70>.

World Health Organization. (2015). The global prevalence of anaemia in 2011. World Health Organization.
<https://apps.who.int/iris/handle/10665/177094>.

Determinants of Iron Supplementation (Fe) Adherence among Pregnant Women at Puskesmas Sugio Lamongan

ORIGINALITY REPORT

19%

SIMILARITY INDEX

15%

INTERNET SOURCES

13%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1	www.ejournal.poltekkes-denpasar.ac.id Internet Source	1%
2	www.annualreviews.org Internet Source	1%
3	Submitted to Institute of Health & Management Pty Ltd Student Paper	1%
4	e-journal.unair.ac.id Internet Source	1%
5	journals.plos.org Internet Source	1%
6	balitsereal.litbang.pertanian.go.id Internet Source	1%
7	Fikadu Waltengus Sendeku, Getnet Gedefaw Azeze, Selamawit Lake Fenta. "Adherence to iron-folic acid supplementation among pregnant women in Ethiopia: a systematic review and meta-analysis", BMC Pregnancy and Childbirth, 2020	1%

8	Pratiwi Cahya Skania, Djaswadi Dasuki, Fitriana Siswi Utami. "The Effect Of Fe Tablet Consumption On Hemoglobin (Hb) Increase In Pregnant Women: A Systematic Literature Review", Jurnal Kebidanan Midwiferia, 2020 Publication	1 %
9	conference.juriskes.com Internet Source	1 %
10	e-journal.sari-mutiara.ac.id Internet Source	1 %
11	ijnpnd.com Internet Source	1 %
12	nrl.northumbria.ac.uk Internet Source	1 %
13	ejournal.almaata.ac.id Internet Source	1 %
14	Submitted to Associatie K.U.Leuven Student Paper	<1 %
15	repository-tnmgrmu.ac.in Internet Source	<1 %
16	Submitted to Adtalem Global Education Student Paper	<1 %
17	Dwiana Kartika Putri. "Hubungan Pengetahuan dan Sikap Ibu Hamil Trimester	<1 %

III Dalam Konsumsi Tablet Fe dengan Terjadinya Anemia Di BPM Mardiani Ilyas Aceh Tahun 2018", Jurnal Midwifery Update (MU), 2019

Publication

18

Muslihatul Jannah, Ni Nengah Arini Murni. "Penggunaan Media Audio Visual Meningkatkan Kepatuhan Konsumsi Tablet Tambah Darah Pada Ibu Hamil", Jurnal Kesehatan Prima, 2019

Publication

<1 %

19

Rosmelia Devia Martha, Sulistyaningsih Sulistyaningsih. "The Administration of Iron Supplements for Anemia Prevention in Pregnant Women: Scoping Review", Jurnal Aisyah : Jurnal Ilmu Kesehatan, 2022

Publication

<1 %

20

Yulia Pratiwi, Tya Safitri. "Kepatuhan Ibu Hamil Dalam Mengonsumsi Tablet Fe (Ferrum) Terhadap Kejadian Anemia Di Desa Langgenharjo Kecamatan Juwana", Lumbung Farmasi: Jurnal Ilmu Kefarmasian, 2021

Publication

<1 %

21

iris.uniroma1.it

Internet Source

<1 %

22

Mohammad Shiddiq Suryadi, Dessy Rindiyanti Harista, Akh. Choirul Abdi. "DESCRIPTION OF COMMUNITY COMPLIANCE IN

<1 %

IMPLEMENTING THE COVID-19 HEALTH
PROTOCOL IN TANGGUMONG VILLAGE,
SAMPANG DISTRICT, REGENCY", Proceeding
of International Conference of Kerta
Cendekia, 2023

Publication

23 Supriyatiningasih Wenang, Arlina Dewi, Ova Emilia, Lidia Febrianti, Joerg Haier. "Key Success Factors in Maternity Waiting Home Implementation: A Qualitative Study in Rural Indonesia", Research Square Platform LLC, 2023 <1 %

Publication

24 Submitted to Universitas Airlangga <1 %

Student Paper

25 docobook.com <1 %

Internet Source

26 ejournal2.litbang.kemkes.go.id <1 %

Internet Source

27 www.scitepress.org <1 %

Internet Source

28 Lacerte, P., M. Pradipasen, P. Temcharoen, N. Imamee, and T. Vorapongsathorn. "Determinants of Adherence to Iron/Folate Supplementation During Pregnancy in Two Provinces in Cambodia", Asia-Pacific Journal of Public Health, 2011. <1 %

29

eurchembull.com

Internet Source

<1 %

30

ji.unbari.ac.id

Internet Source

<1 %

31

repository.unimugo.ac.id

Internet Source

<1 %

32

www.midwifery.iocspublisher.org

Internet Source

<1 %

33

Aviva I Rappaport, Kyly C Whitfield, Gwen E Chapman, Rickey Y Yada et al. "Randomized controlled trial assessing the efficacy of a reusable fish-shaped iron ingot to increase hemoglobin concentration in anemic, rural Cambodian women", *The American Journal of Clinical Nutrition*, 2017

Publication

<1 %

34

Shaymaa Abdullateef Alfadhul, Shiamaa Mohammed Ali Kadhum. "The significance of awareness about iron deficiency anaemia in its prevention among Iraqi pregnant women attending primary healthcare centres", *Family Medicine & Primary Care Review*, 2023

Publication

<1 %

35

aisyah.journalpress.id

Internet Source

<1 %

36	irjpms.com Internet Source	<1 %
37	journal.poltekkes-mks.ac.id Internet Source	<1 %
38	jurnal.htp.ac.id Internet Source	<1 %
39	midwifery.iocspublisher.org Internet Source	<1 %
40	www.thieme-connect.com Internet Source	<1 %
41	Abebe Mekonnen, Walleign Alemnew, Zegeye Abebe, Getu Debalkie Demissie. "Adherence to Iron with Folic Acid Supplementation Among Pregnant Women Attending Antenatal Care in Public Health Centers in Simada District, Northwest Ethiopia: Using Health Belief Model Perspective", Patient Preference and Adherence, 2021 Publication	<1 %
42	Ali Sungkar, Saptawati Bardosono, Rima Irwinda, Nurul R. M. Manikam et al. "A Life Course Approach to the Prevention of Iron Deficiency Anemia in Indonesia", Nutrients, 2022 Publication	<1 %

43

Djibril M Ba, Paddy Ssentongo, Kristen H Kjerulff, Muzi Na, Guodong Liu, Xiang Gao, Ping Du. "Adherence to Iron Supplementation in 22 Sub-Saharan African Countries and Associated Factors among Pregnant Women: A Large Population-Based Study", Current Developments in Nutrition, 2019

Publication

<1 %

44

Takele Gezahegn Demie, Getachew Tilahun Gessese, Berhanu Teshome Woldeamanuel, Tolesa Diriba Biratu, Simegnew Handebo. "Adherence to iron supplement intake during pregnancy and associated factors in Ethiopia: Further analysis of a national population-based study", Food Science & Nutrition, 2023

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On